

TWISTING OF QUANTUM SPACES AND TWISTED COHOM OBJECTS

SERGIO D. GRILLO

ABSTRACT. Twisting process for quantum linear spaces is defined. It consists in a particular kind of globally defined deformations on finitely generated algebras. Given a quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$, a multiplicative cosimplicial quasicomplex $\mathbf{C}^\bullet[\mathbf{A}_1]$ in the category Grp is associated to \mathbf{A}_1 , in such a way that for every $n \in \mathbb{N}$ a subclass of linear automorphisms of $\mathbf{A}^{\otimes n}$ is obtained from the groups $\mathbf{C}^n[\mathbf{A}_1]$. Among the elements of this subclass, the counital 2-cocycles are those which define the twist transformations. In these terms, the twisted internal coHom objects $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$, constructed in [6], can be described as twisting of the proper coHom objects $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$. Moreover, twisted tensor products $\mathcal{A} \circ_\tau \mathcal{B}$, in terms of which above objects were built up, can be seen as particular 2-cocycle twisting of $\mathcal{A} \circ \mathcal{B}$, enabling us to generalize the mentioned construction. The quasicomplexes $\mathbf{C}^\bullet[\mathbf{V}]$ are further studied, showing for instance that, when \mathbf{V} is a coalgebra the quasicomplexes related to Drinfeld twisting, corresponding to bialgebras generated by \mathbf{V} , are subobjects of $\mathbf{C}^\bullet[\mathbf{V}]$.

INTRODUCTION

In a previous paper [6], combining the ideas of twisted tensor products (TTP) [1] and internal coHom objects, we have built up on the monoidal category $(\text{CA}, \circ, \mathcal{K})$ of *conic algebras* or *conic quantum spaces*, i.e. finitely generated graded algebras (which constitutes a full subcategory of the -general- quantum linear spaces [12][13]), functions $(\mathcal{B}, \mathcal{A}) \mapsto \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \in \text{CA}$, one for each collection of automorphisms $\Upsilon = \{\sigma_{\mathcal{A}} : \mathcal{A} \simeq \mathcal{A}\}_{\mathcal{A} \in \text{CA}}$, defining CA^{op} -based categories distinct from the one related to its proper internal coHom objects $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] = \mathcal{B} \triangleright \mathcal{A}$ (in the sense that, in general, $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \not\cong \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$). The opposite objects

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]^{op} \doteq \underline{\text{Hom}}^\Upsilon[\mathcal{B}^{op}, \mathcal{A}^{op}] \in \text{CA}^{op}$$

represent ‘spaces of morphisms’ whose coordinate rings, given precisely by the algebras $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$, do not commute with the ones of their respective domain \mathcal{B}^{op} , in the sense that they give rise to *twisted coevaluation* arrows $\mathcal{A} \rightarrow \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \circ_\tau \mathcal{B}$, where \circ_τ is a TTP constructed in terms of $\sigma_{\mathcal{A}}, \sigma_{\mathcal{B}} \in \Upsilon$. This is why we have called them *twisted coHom* objects.

We have seen in that paper the objects $\underline{\text{end}}^\Upsilon[\mathcal{A}] = \underline{\text{hom}}^\Upsilon[\mathcal{A}, \mathcal{A}]$, which are endowed with a bialgebra structure, are counital 2-cocycle twisting [4] of the coEnd objects $\underline{\text{end}}[\mathcal{A}] = \mathcal{A} \triangleright \mathcal{A}$. In other words, those objects are related by a (non formal) bialgebra *deformation*. However, we have not been able to make an analogous claim relating the objects $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ and $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$ for $\mathcal{B} \neq \mathcal{A}$ yet. To this end, we will develop in this paper a suitable kind of globally defined deformations of quantum spaces, which we shall call *twisting of quantum spaces*, in such a way that a multiplicative cosimplicial quasicomplex structure \mathbf{C}^\bullet can be related to them, being 2-cocycles the *well-behaved* deformations. We shall see the objects $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ are twisting by 2-cocycles of $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$. In particular, for the coEnd objects, those transformations define the previously cited 2-cocycle bialgebra twisting.

The main aim of our work is to construct and analyze the mentioned cosimplicial quasicomplex structure and related algebra deformation process. We invoke the concept of twisted coHom objects just because they were our main motivation to this paper.

Although our twist transformation will be defined on the hole class of quantum spaces, i.e. the category FGA of finitely generated algebras, they are mainly designed to be applied on the conic ones. This is why, among other things, the first chapter is dedicated to them. Thus, in §1 we make a brief review about conic quantum spaces, recalling their monoidal structures and related internal (co)Hom objects. For a more extended treatment, reference [6] can be consulted.

In order to introduce the twisting process in a (quasi)cohomological framework, in §2 we endow every linear space \mathbf{V} (or every tensor algebra \mathbf{V}^{\otimes}) with a multiplicative cosimplicial quasicomplex structure $(\mathbf{C}^\bullet[\mathbf{V}], \partial)$. Recall that formal deformations [5] of an associative algebra \mathbf{A} are controlled by the Hochschild complex [8] of \mathbf{A} . Given a pair $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ and a (counital) 2-cocycle ψ inside certain subclass of $\mathbf{C}^2[\mathbf{A}_1]$, the *admissible* 2-cochains, we define in §3 a deformation of \mathcal{A} as a new quantum space \mathcal{A}_ψ , the *twisting* of \mathcal{A} by ψ . The mentioned subclass defines a subgroup of linear automorphisms of $\mathbf{A} \otimes \mathbf{A}$. Deformed quantum spaces \mathcal{A}_ψ and

Date: May 2002.

1991 *Mathematics Subject Classification*. Primary 46L65, 46L52; Secondary 18G50.

\mathcal{A}_ψ are isomorphic *iff* the cocycles ψ and φ are cohomologous through some admissible 1-cochain. We show that twist transformation can be composed and have inverse. Moreover, an equivalence relation between quantum spaces can be defined, namely: \mathcal{A} and \mathcal{B} are *twist or gauge related* when there exists ψ such that $\mathcal{A}_\psi \simeq \mathcal{B}$ (as in the case of bialgebra twisting). We also show the twisting of bialgebras in the category FGA are particular twisting of quantum spaces.

Finally, we investigate in §4 the behavior of twist transformations in relation to certain functorial structures in FGA. We see, for example, that products $\mathcal{A}_\psi \circ \mathcal{B}_\varphi$ are twisting of products $\mathcal{A} \circ \mathcal{B}$. Analogous claim is also true, when restricted to particular kind of twist transformations, for functors $!$, \bullet , and products \odot , \triangleleft , \triangleright and \diamond defined in [6]. This study lead us to a better understanding of the construction of twisted coHom objects and, in turn, a way to generalize it.

We often adopt definitions and notation extracted form Mac Lane's book [10]. \mathbb{k} indicates some of the numeric fields, \mathbb{R} or \mathbb{C} . The usual tensor product on $\mathbb{k}\text{-Alg} \equiv \text{Alg}$ and $\text{Vct}_{\mathbb{k}} \equiv \text{Vct}$ (the categories of unital associative \mathbb{k} -algebras and of \mathbb{k} -vector spaces, respectively) is denoted by \otimes . Grp denotes the category of groups and their homomorphisms.

1. CONIC QUANTUM SPACES

In this chapter we recall definition, functorial structures and internal coHom objects (standard and twisted ones) of a subclass of quantum spaces that we have called *conic quantum spaces* [6], introducing the needed notation for the rest of the paper. We suppose the reader is familiar with the concepts of quadratic and general quantum spaces [12][13].

Conic quantum spaces define a category CA such that the full inclusions $\text{QA} \subset \text{CA} \subset \text{FGA}$ holds, being FGA and QA the categories of finitely generated algebras and of quadratic algebras, respectively. Remember that FGA is formed out by pairs $(\mathbf{A}_1, \mathbf{A})$ where \mathbf{A} is a unital algebra generated by a finite linear subspace $\mathbf{A}_1 \subset \mathbf{A}$, and its morphisms are algebra morphisms $\mathbf{A} \rightarrow \mathbf{B}$ such that restricted to $\mathbf{A}_1 \subset \mathbf{A}$ give linear maps $\mathbf{A}_1 \rightarrow \mathbf{B}_1$ ($\subset \mathbf{B}$). The objects of CA are pairs $\mathcal{A} \doteq (\mathbf{A}_1, \mathbf{A}) \in \text{FGA}$ such that \mathbf{A} is a graded algebra with grading given by

$$\mathbf{A} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{A}_n; \quad \mathbf{A}_n = \Pi(\mathbf{A}_1^{\otimes n}); \quad \mathbb{N}_0 = \mathbb{N} \cup \{0\},$$

where $\Pi = \Pi_{\mathbf{A}} : \mathbf{A}_1^{\otimes} \rightarrow \mathbf{A}$ is the canonical epimorphism of algebras such that restricted to \mathbf{A}_1 gives the inclusion $\mathbf{A}_1 \hookrightarrow \mathbf{A}$. We are denoting by \mathbf{A}_1^{\otimes} the tensor algebra of \mathbf{A}_1 , i.e. $\mathbf{A}_1^{\otimes} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{A}_1^{\otimes n}$. In the general case the related algebras are just filtered, more precisely

$$(1.1) \quad \mathbf{A} = \bigcup_{n \in \mathbb{N}_0} \mathbf{F}_n; \quad \mathbf{F}_n = \Pi\left(\bigoplus_{i=0}^n \mathbf{A}_1^{\otimes i}\right).$$

Alternatively, the objects of CA can be described as those pairs \mathcal{A} in FGA such that $\ker \Pi$ is a graded bilateral ideal $\ker \Pi_{\mathbf{A}} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n$ with $\mathbf{I}_n \subset \mathbf{A}_1^{\otimes n}$. In these terms, given another pair $(\mathbf{B}_1, \mathbf{B})$ with $\ker \Pi_{\mathbf{B}} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{J}_n$, the arrows $(\mathbf{A}_1, \mathbf{A}) \rightarrow (\mathbf{B}_1, \mathbf{B})$ in CA are characterized by linear maps $\alpha_1 : \mathbf{A}_1 \rightarrow \mathbf{B}_1$ such that $\alpha_1^{\otimes n}(\mathbf{I}_n) \subset \mathbf{J}_n$, being

$$\alpha_1^{\otimes} = \bigoplus_{n \in \mathbb{N}_0} \alpha_1^{\otimes n} : \mathbf{A}_1^{\otimes} \rightarrow \mathbf{B}_1^{\otimes}$$

the unique extension of α_1 to \mathbf{A}_1^{\otimes} as a morphism of algebras. The algebra map $\alpha : \mathbf{A} \rightarrow \mathbf{B}$ that defines any morphism in FGA is the unique algebra homomorphism such that $\alpha \Pi_{\mathbf{A}} = \Pi_{\mathbf{B}} \alpha_1^{\otimes}$. In particular, $\alpha_1 = \alpha|_{\mathbf{A}_1}$. Returning to the ideal of a conic quantum space, since the restriction of $\Pi_{\mathbf{A}}$ to \mathbf{A}_1 is the identity, we always have $\mathbf{I}_1 = \{0\}$. In addition, $\mathbf{I}_0 \neq \{0\}$ *iff* $\mathbf{A} = \{0\}$. Thus, generically $\ker \Pi_{\mathbf{A}} = \bigoplus_{n \geq 2} \mathbf{I}_n$, unless $\mathbf{A} = \{0\}$.

Examples of conic quantum spaces are, beside quadratics, those with $\ker \Pi = I[\mathbf{Y}]$, where $\mathbf{Y} \subset \mathbf{A}_1^{\otimes m}$ for some $m > 2$, indicating by $I[\mathbf{X}]$ the bilateral ideal generated algebraically by $\mathbf{X} \subset \mathbf{A}_1^{\otimes}$; i.e. $I[\mathbf{X}] = \mathbf{A}_1^{\otimes} \otimes \mathbf{X} \otimes \mathbf{A}_1^{\otimes}$. We have called them *m-th quantum spaces*, and denoted CA^m the full subcategory of FGA which has these pairs as objects. Thus $\text{QA} = \text{CA}^2$.

We recall that from the geometric point of view, the quantum spaces actually are the opposite objects \mathcal{A}^{op} to the pairs $\mathcal{A} = (\mathbf{A}_1, \mathbf{A}) \in \text{FGA}$, being the latter their (generically) *noncommutative coordinate rings*. That is to say FGA^{op} , which we indicate QLS, is the category of quantum spaces, and FGA the category of their coordinate rings. Nevertheless, since the duality between QLS and FGA, for quantum spaces we understand the objects of each one of these categories. Categories CA and CA^m give rise to full subcategories of QLS.

1.1. **Functorial structures and internal coHom objects.** The bifunctor \circ defines an (strict and symmetric) monoidal structure with unit $\mathcal{K} = (\mathbb{k}, \mathbb{k}^\otimes)$ in CA. Recall that $\mathcal{A} \circ \mathcal{B} = (\mathbf{A}_1 \otimes \mathbf{B}_1, \mathbf{A} \circ \mathbf{B})$, being $\mathbf{A} \circ \mathbf{B}$ the subalgebra of $\mathbf{A} \otimes \mathbf{B}$ generated by $\mathbf{A}_1 \otimes \mathbf{B}_1$, and that the unit for \circ is $\mathcal{I} \doteq (\mathbb{k}, \mathbb{k})$ in FGA. If $\mathcal{A}, \mathcal{B} \in \text{CA}$, the algebra $\mathbf{A} \circ \mathbf{B}$ effectively defines an object of CA due to the kernel of $\Pi_\circ : [\mathbf{A}_1 \otimes \mathbf{B}_1]^\otimes \rightarrow \mathbf{A} \circ \mathbf{B}$, using the (canonical) homogeneous isomorphism of graded algebras

$$(1.2) \quad [\mathbf{A}_1 \otimes \mathbf{B}_1]^\otimes \simeq [\mathbf{A}_1 \otimes \mathbf{B}_1]^{\hat{\otimes}} \doteq \bigoplus_{n \in \mathbb{N}_0} (\mathbf{A}_1^{\otimes n} \otimes \mathbf{B}_1^{\otimes n}),$$

being the latter the subalgebra of $\mathbf{A}_1^\otimes \otimes \mathbf{B}_1^\otimes$ generated by $\mathbf{A}_1 \otimes \mathbf{B}_1$, is a graded bilateral ideal isomorphic to

$$\bigoplus_{n \in \mathbb{N}_0} (\mathbf{I}_n \otimes \mathbf{B}_1^{\otimes n} + \mathbf{A}_1^{\otimes n} \otimes \mathbf{J}_n) \left(\subset [\mathbf{A}_1 \otimes \mathbf{B}_1]^{\hat{\otimes}} \right).$$

Note that $[\mathbf{A}_1 \otimes \mathbf{B}_1]^{\hat{\otimes}} = \mathbf{A}_1^\otimes \circ \mathbf{B}_1^\otimes$. We frequently identify the latter algebra and $[\mathbf{A}_1 \otimes \mathbf{B}_1]^\otimes$. In particular, Π_\circ will also be understood as a map $\mathbf{A}_1^\otimes \circ \mathbf{B}_1^\otimes \rightarrow \mathbf{A} \circ \mathbf{B}$.

For quadratic algebras there is another monoidal structure \bullet and a functor $!$ such that

$$(1.3) \quad \mathcal{A}^! \simeq \mathcal{A}, \quad (\mathcal{A} \circ \mathcal{B})^! \simeq \mathcal{A}^! \bullet \mathcal{B}^!, \quad (\mathcal{A} \bullet \mathcal{B})^! \simeq \mathcal{A}^! \circ \mathcal{B}^!, \quad \mathcal{K}^! \simeq \mathcal{U},$$

being $\mathcal{U} = (\mathbb{k}, \mathbf{U})$, $\mathbf{U} = \mathbb{k}[e]/I[e^2]$, a unit object for \bullet . Analogous functors to \bullet and $!$ can be given in CA, namely

$$\mathbf{A} \circ \mathbf{B} \doteq \mathbf{A}_1^\otimes \circ \mathbf{B}_1^\otimes / \bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n \otimes \mathbf{J}_n$$

and

$$(1.4) \quad \mathbf{A}^! \doteq \mathbf{A}_1^{*\otimes} / \mathbf{I}^\dagger; \quad \mathbf{I}^\dagger \doteq \bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n^\dagger \doteq I \left[\bigoplus_{n \geq 2} \mathbf{I}_n^\perp \right],$$

with $\mathbf{I}_n^\perp = \{x \in \mathbf{A}_1^{*\otimes n} : \langle x, y \rangle = 0, \forall y \in \mathbf{I}_n\}$, for $n \geq 2$. We shall take $\mathbf{I}_{0,1}^\perp = \{0\}$. A unit element for (CA, \circ) is \mathcal{U} as for QA, but the restriction of \circ to QA does not coincide with the functor \bullet . On the other hand, the functor $!$ does coincide with the corresponding to the quadratic cases, although $!^2 \not\cong id_{\text{CA}}$. The preserved properties are, with respect to (1.3),

$$(\mathcal{A} \circ \mathcal{B})^! \simeq \mathcal{A}^! \circ \mathcal{B}^!, \quad \mathcal{K}^! \simeq \mathcal{U}, \quad \mathcal{K} \simeq \mathcal{U}^!,$$

while $(\mathcal{A} \circ \mathcal{B})^! \not\cong \mathcal{A}^! \circ \mathcal{B}^!$ and $\mathcal{A}^{!!} \not\cong \mathcal{A}$. Some kind of covariant mixing of \circ and $!$ can be defined, namely

$$\triangleright : \text{CA}^{op} \times \text{CA} \rightarrow \text{CA}, \quad \triangleleft : \text{CA} \times \text{CA}^{op} \rightarrow \text{CA},$$

$$\diamond : \text{CA}^{op} \times \text{CA}^{op} \rightarrow \text{CA},$$

satisfying

$$(1.5) \quad \triangleright = \bullet (! \times id), \quad \triangleleft = \bullet (id \times !), \quad \diamond = \bullet (! \times !)$$

when restricted to QA. They are given on objects by (identifying each object with its opposite)

$$\mathbf{A} \triangleright \mathbf{B} \doteq \mathbf{A}_1^{*\otimes} \circ \mathbf{B}_1^\otimes / I \left[\bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n^\perp \otimes \mathbf{J}_n \right],$$

$$\mathbf{A} \triangleleft \mathbf{B} \doteq \mathbf{A}_1^\otimes \circ \mathbf{B}_1^{*\otimes} / I \left[\bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n \otimes \mathbf{J}_n^\perp \right],$$

$$\mathbf{A} \diamond \mathbf{B} \doteq \mathbf{A}_1^{*\otimes} \circ \mathbf{B}_1^{*\otimes} / I \left[\bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n^\perp \otimes \mathbf{J}_n^\perp \right].$$

The functors \triangleright and \triangleleft have \mathcal{K} as left and as a right unit, respectively, in the sense that $\mathcal{K} \triangleright \mathcal{A} \simeq \mathcal{A}$ and $\mathcal{A} \triangleleft \mathcal{K} \simeq \mathcal{A}$ for any \mathcal{A} in CA (on the other hand, $\mathcal{A} \triangleright \mathcal{U} \simeq \mathcal{A}^!$, $\mathcal{U} \triangleleft \mathcal{A} \simeq \mathcal{A}^!$ and $\mathcal{K} \diamond \mathcal{A} \simeq \mathcal{A} \diamond \mathcal{K} \simeq \mathcal{A}^!$).

While the internal coHom objects of QA are given by $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] = \mathcal{B}^! \bullet \mathcal{A}$, for CA, as it might be expected from Eq. (1.5), we have the following result.

Theorem 1. *The category $(\text{CA}, \circ, \mathcal{K})$ has **internal coHom objects** given by $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] = \mathcal{B} \triangleright \mathcal{A}$ with coevaluation arrow $\mathcal{A} \rightarrow (\mathcal{B} \triangleright \mathcal{A}) \circ \mathcal{B}$ defined by the map $a_i \mapsto (b^j \otimes a_i) \otimes b_j$ (sum over repeated indices is understood), where $\{a_i\}$, $\{b_j\}$ and $\{b^j\}$ are basis of \mathbf{A}_1 , \mathbf{B}_1 and \mathbf{B}_1^* , respectively. ■*

The proof is given in [6].

1.2. The twisted internal coHom objects. An internal coHom object in FGA is an initial object of the comma category $(\mathcal{A} \downarrow \text{FGA} \circ \mathcal{B})$. The objects of each $(\mathcal{A} \downarrow \text{FGA} \circ \mathcal{B})$, *diagrams* in the Manin terminology, are pairs $\langle \varphi, \mathcal{H} \rangle_{\mathcal{A}, \mathcal{B}} = \langle \varphi, \mathcal{H} \rangle$, with $\mathcal{H} \in \text{FGA}$ and φ a morphism $\mathcal{A} \rightarrow \mathcal{H} \circ \mathcal{B}$; and its arrows $\langle \varphi, \mathcal{H} \rangle \rightarrow \langle \varphi', \mathcal{H}' \rangle$ are given by morphisms $\alpha : \mathcal{H} \rightarrow \mathcal{H}'$ satisfying $\varphi' = (\alpha \circ I_{\mathcal{B}}) \varphi$. For every such category $\langle \varphi, \mathcal{H} \rangle \mapsto \mathcal{H}$ defines an embedding $\mathbb{U} : (\mathcal{A} \downarrow \text{FGA} \circ \mathcal{B}) \hookrightarrow \text{FGA}$. The disjoint union of the family $\{(\mathcal{A} \downarrow \text{FGA} \circ \mathcal{B})\}_{\mathcal{A}, \mathcal{B} \in \text{FGA}}$, namely FGA° , has a semigroupoid structure given by the partial product functor $\widehat{\circ}$. \mathbb{U} extends to an obvious embedding $\text{FGA}^\circ \hookrightarrow \text{FGA}$, which we shall also call \mathbb{U} , and that satisfies

$$(1.6) \quad \mathbb{U} \widehat{\circ} = \circ (\mathbb{U} \times \mathbb{U}) \quad \text{and} \quad \mathbb{U} \langle \ell_{\mathcal{A}}, \mathcal{I} \rangle = \mathcal{I}.$$

So $\mathbb{U} : \text{FGA}^\circ \hookrightarrow \text{FGA}$ is an embedding of categories with unital associative partial products. The same is true for CA and every CA^m , but changing \mathcal{I} by \mathcal{K} in Eq. (1.6). We have seen in [6] that it is possible to arrive at the notions of cocomposition and coidentity from the semigroupoid structure in FGA° , the initiality of each $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \in (\mathcal{A} \downarrow \text{FGA} \circ \mathcal{B})$, and the existence of a functor \mathbb{U} satisfying Eq. (1.6). Then, the function $(\mathcal{B}, \mathcal{A}) \mapsto \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$, together with the corresponding cocomposition and coidentity arrows, defines an FGA-cobased (or dually FGA^{op} -based) category with objects also in FGA. Such a cobased category has in addition the usual notion of coevaluation $\mathcal{A} \rightarrow \underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \circ \mathcal{B}$.

Based on these ideas, we construct in that paper a family of categories $\Upsilon^{\mathcal{A}, \mathcal{B}}$ with initial objects, denoted $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$, and with a family of embedding $\Upsilon^{\mathcal{A}, \mathcal{B}} \hookrightarrow \text{FGA}$. The objects of each $\Upsilon^{\mathcal{A}, \mathcal{B}}$ are essentially arrows $\mathcal{A} \rightarrow \mathcal{H} \circ_\tau \mathcal{B}$, where \circ_τ denotes a twisted tensor product between \mathcal{H} and \mathcal{B} . Each category is defined by a linear isomorphism $\widehat{\tau}_{\mathcal{A}, \mathcal{B}} : \mathbf{B}_1 \otimes \mathbf{B}_1^* \otimes \mathbf{A}_1 \simeq \mathbf{B}_1 \otimes \mathbf{B}_1^* \otimes \mathbf{A}_1$ from which the twisted products \circ_τ are built up. Restricting \mathcal{A}, \mathcal{B} to CA and considering $\widehat{\tau}_{\mathcal{A}, \mathcal{B}} = \text{id} \otimes \sigma_{\mathcal{B}}^{*-1} \otimes \sigma_{\mathcal{A}}$, the disjoint union Υ^\cdot of these categories has a semigroupoid structure such that an equation like (1.6) holds for the extended functor $\Upsilon^\cdot \hookrightarrow \text{CA}$. Thus, $(\mathcal{B}, \mathcal{A}) \mapsto \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ gives rise to an CA-cobased category with the additional notion of coevaluation $\mathcal{A} \rightarrow \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \circ_\tau \mathcal{B}$. We will not give the details of this construction, referring the interested reader to [6]. Nevertheless, in the rest of this section we review the main properties of the objects $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ and its relationship to the proper coHom objects $\mathcal{B} \triangleright \mathcal{A}$ of CA.

Let $\{\sigma_{\mathcal{A}} : \mathbf{A}_1 \simeq \mathbf{A}_1\}_{\mathcal{A} \in \text{CA}}$ be a collection defining Υ^\cdot , such that each map $\sigma_{\mathcal{A}}$ can be extended to an automorphism $\mathcal{A} \simeq \mathcal{A}$. Consider a couple \mathcal{A} and \mathcal{B} of quantum spaces given by algebras $\mathbf{A} \simeq \mathbf{A}_1^\otimes / \mathbf{I}$ and $\mathbf{B} \simeq \mathbf{B}_1^\otimes / \mathbf{J}$, being \mathbf{I} and \mathbf{J} the ideals linearly generated by

$$(1.7) \quad R_{\lambda_n}^{k_1 \dots k_n} a_{k_1} \dots a_{k_n} \in \mathbf{I}_n, \quad S_{\mu_n}^{k_1 \dots k_n} b_{k_1} \dots b_{k_n} \in \mathbf{J}_n,$$

with λ_n and μ_n in index sets Λ_n and Φ_n , respectively. Writing $\phi \doteq \sigma_{\mathcal{A}}$ and $\rho \doteq \sigma_{\mathcal{B}}^{*-1}$, the initial objects of each category $\Upsilon^{\mathcal{A}, \mathcal{B}}$ can be defined by a conic quantum space

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] = \mathcal{B}^\Upsilon \triangleright \mathcal{A}^\Upsilon, \quad \mathcal{A}^\Upsilon \doteq (\mathbf{A}_1, \mathbf{A}_1^\otimes / \mathbf{I}_\sigma), \quad \mathcal{B}^\Upsilon \doteq (\mathbf{B}_1, \mathbf{B}_1^\otimes / \mathbf{J}_\sigma),$$

with \mathbf{I}_σ and \mathbf{J}_σ linearly generated by

$$\left\{ \left\{ \left\{ R_{\lambda_n}^{k_1 \dots k_n} a_{k_1} \dots a_{k_n} \right\}_{\lambda_n \in \Lambda_n} \right\}_{n \in \mathbb{N}_0}, \left\{ \left\{ \left\{ S_{\mu_n}^{k_1 \dots k_n} b_{k_1} \dots b_{k_n} \right\}_{\mu_n \in \Phi_n} \right\}_{n \in \mathbb{N}_0} \right\},$$

where

$$(1.8) \quad {}_\sigma R_{\lambda_n}^{k_1 \dots k_n} \doteq R_{\lambda_n}^{k_1 j_2 \dots j_n} \phi_{j_2}^{k_2} (\phi^2)_{j_3}^{k_3} \dots (\phi^{n-1})_{j_n}^{k_n},$$

$$(1.9) \quad {}_\sigma S_{\mu_n}^{k_1 \dots k_n} \doteq S_{\mu_n}^{k_1 j_2 \dots j_n} (\rho^{-1})_{j_2}^{k_2} (\rho^{-2})_{j_3}^{k_3} \dots (\rho^{1-n})_{j_n}^{k_n}.$$

On the other hand, if each $\mathbf{J}_n^\perp \subset \mathbf{B}_1^{*\otimes n}$ is the span of

$$(1.10) \quad \left\{ b^{k_1} \dots b^{k_n} (S^\perp)_{k_1 \dots k_n}^{\omega_n} \right\}_{\omega_n \in \Omega_n},$$

$(\mathbf{J}_\sigma)_n^\perp$ is spanned by the set

$$(1.11) \quad \left\{ b^{k_1} b^{k_2} \dots b^{k_n} ({}_\sigma S^\perp)_{k_1 \dots k_n}^{\omega_n} \right\}_{\omega_n \in \Omega_n},$$

with

$$(1.12) \quad ({}_\sigma S^\perp)_{k_1 \dots k_n}^{\omega_n} \doteq \rho_{k_2}^{j_2} (\rho^2)_{k_3}^{j_3} \dots (\rho^{n-1})_{k_n}^{j_n} (S^\perp)_{k_1 j_2 \dots j_n}^{\omega_n}.$$

Therefore, $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ will be the algebra generated by $z_i^j = b^j \otimes a_i$ and quotient by the ideal algebraically generated by

$$\left\{ \left\{ \left\{ {}_\sigma R_{\lambda_n}^{k_1 \dots k_n} z_{k_1}^{j_1} \cdot z_{k_2}^{j_2} \dots z_{k_n}^{j_n} ({}_\sigma S^\perp)_{j_1 \dots j_n}^{\omega_n} \right\}_{\omega_n \in \Omega_n} \right\}_{n \in \mathbb{N}_0} \right\}.$$

From the semigroupoid structure of Υ and the embedding $\Upsilon^{\mathcal{A}, \mathcal{B}} \hookrightarrow \text{CA}$, there exist arrows

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \rightarrow \underline{\text{hom}}^\Upsilon[\mathcal{C}, \mathcal{A}] \circ \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{C}], \quad \underline{\text{end}}^\Upsilon[\mathcal{A}] \rightarrow \mathcal{K},$$

giving us the notions of cocomposition and coidentity we have just mentioned. Of course, these arrows define a bialgebra structure on $\underline{\text{end}}^\Upsilon[\mathcal{A}]$. Moreover, there exists a counital 2-cocycle $\chi_{\mathcal{A}} = \chi$, such that $\underline{\text{end}}^\Upsilon[\mathcal{A}] \simeq (\mathcal{A} \triangleright \mathcal{A})_\chi$. That means $\underline{\text{end}}^\Upsilon[\mathcal{A}]$ is a twisting by χ of $\underline{\text{end}}[\mathcal{A}]$ [4]. Moreover, from the actions $\mathcal{A} \rightarrow \underline{\text{end}}[\mathcal{A}] \circ \mathcal{A}$ (the standard coevaluation maps for CA), the above twisting can be translated to each \mathcal{A} defining a new quantum space \mathcal{A}_χ isomorphic to \mathcal{A}^Υ (c.f. [11], page 54). Finally,

$$\underline{\text{end}}^\Upsilon[\mathcal{A}] \simeq \mathcal{A}_\chi \triangleright \mathcal{A}_\chi = \underline{\text{end}}[\mathcal{A}_\chi] \simeq \underline{\text{end}}[\mathcal{A}]_\chi.$$

This paper was mainly motivated to make valid the last equation for every coHom object. Note that for $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$, with $\mathcal{B} \neq \mathcal{A}$, only a part of this equation is valid, i.e.

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \simeq \mathcal{B}_\chi \triangleright \mathcal{A}_\chi = \underline{\text{hom}}[\mathcal{B}_\chi, \mathcal{A}_\chi],$$

since the last part has non sense for a quantum space which has not a bialgebra structure. To do that we shall define along the next chapters a twisting process for all quantum spaces.

2. COSIMPLICIAL QUASICOMPLEXES FOR TENSOR ALGEBRAS

Let us consider a (general) quantum space $(\mathbf{A}_1, \mathbf{A})$ with related algebra structure (m, η) . Any linear endomorphism $\Xi \in \text{End}_{\text{Vct}}[\mathbf{A} \otimes \mathbf{A}]$ defines a new product¹ $m_\Xi \doteq m \Xi$ on \mathbf{A} , which can be called *twisting or deformation of m by Ξ* . Let us suppose Ξ is such that m_Ξ is associative. If in addition Ξ satisfies

$$(2.1) \quad \Xi(\eta \otimes I) = \eta \otimes I \quad \text{and} \quad \Xi(I \otimes \eta) = I \otimes \eta,$$

then η is also a unit for m_Ξ and we have a new algebra structure (m_Ξ, η) on \mathbf{A} , namely the *twisting or deformation \mathbf{A}_Ξ of \mathbf{A} by Ξ* . In general, the pair $(\mathbf{A}_1, \mathbf{A}_\Xi)$ is not a quantum space, because the vector space generated algebraically by \mathbf{A}_1 through the product m_Ξ is not all of $\mathbf{A} = \mathbf{A}_\Xi$ (the equality is for the underlying vector spaces). Just take $\Xi \equiv 0$. Things change when we restrict ourself to the set $\text{Aut}_{\text{Vct}}[\mathbf{A} \otimes \mathbf{A}]$, as we shall see in §3.1. On the other hand, if $(\mathbf{A}_1, \mathbf{A})$ is a conic quantum space with $\mathbf{A} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{A}_n$, we can ask Ξ to be a *bihomogeneous* of degree zero, i.e. $\Xi(\mathbf{A}_r \otimes \mathbf{A}_s) \subset \mathbf{A}_r \otimes \mathbf{A}_s$, $\forall r, s \in \mathbb{N}_0$. In such a case m_Ξ gives rise clearly to the same gradation as m . Thus, our objects of interest are homogeneous linear automorphisms satisfying (2.1).

But, what kind of condition(s) must be imposed on Ξ to insure m_Ξ is associative (when m is). We shall show immediately that there is a cosimplicial quasicomplex structure related to every tensor algebra of a given vector space. Since every linear map in \mathbf{A} can be defined by one in \mathbf{A}_1^\otimes through the epimorphism $\mathbf{A}_1^\otimes \rightarrow \mathbf{A}$, we will study the maps on $\mathbf{A} \otimes \mathbf{A}$ in terms of those on $\mathbf{A}_1^\otimes \otimes \mathbf{A}_1^\otimes$ and the mentioned quasicomplex structure. As one might expect, 2-cocycles will define maps whose related twisted products are associative.

Our construction of the cosimplicial object and the associated cochain quasicomplex \mathbf{C}^\bullet was inspired by the one developed in [11] and related to twisting of bialgebras,² which we shall call \mathbf{G}^\bullet . At the end of this chapter the relationship between these quasicomplexes will be analyzed.

2.1. The cosimplicial object. Consider $\mathbf{V} \in \text{Vct}$ and its associated tensor algebra $\mathbf{V}^\otimes = \bigoplus_{n \in \mathbb{N}_0} \mathbf{V}^{\otimes n}$, $\mathbf{V}^{\otimes 0} = \mathbf{k}$. We shall call

$$\mathbf{C}^n[\mathbf{V}] \subset \text{Aut}_{\text{Vct}}[(\mathbf{V}^\otimes)^{\otimes n}], \quad n \in \mathbb{N}_0,$$

the subgroup of *n-homogeneous* of degree zero, or simply homogeneous, linear automorphisms of $(\mathbf{V}^\otimes)^{\otimes n}$, i.e. those linear maps $\psi : (\mathbf{V}^\otimes)^{\otimes n} \simeq (\mathbf{V}^\otimes)^{\otimes n}$ such that the equalities of sets

$$(2.2) \quad \psi(\mathbf{V}^{\otimes r_1} \otimes \dots \otimes \mathbf{V}^{\otimes r_n}) = \mathbf{V}^{\otimes r_1} \otimes \dots \otimes \mathbf{V}^{\otimes r_n}$$

hold for every $r_k \in \mathbb{N}_0$ and $k = 1 \dots n$. In the finite dimensional case, such conditions are equivalent to the inclusions

$$\psi(\mathbf{V}^{\otimes r_1} \otimes \dots \otimes \mathbf{V}^{\otimes r_n}) \subset \mathbf{V}^{\otimes r_1} \otimes \dots \otimes \mathbf{V}^{\otimes r_n}.$$

The unit element of $\mathbf{C}^n[\mathbf{V}]$ is $\mathbb{I}^{\otimes n}$, being \mathbb{I} the identity map of $\text{Aut}_{\text{Vct}}[\mathbf{V}^\otimes]$. In terms of *n-fold multi-index* $R = (r_1, \dots, r_n) \in \mathbb{N}_0^{\times n}$, we can write

$$(2.3) \quad (\mathbf{V}^\otimes)^{\otimes n} = \bigoplus_{R \in \mathbb{N}_0^{\times n}} \mathbf{V}^{\otimes R}; \quad \mathbf{V}^{\otimes R} \doteq \mathbf{V}^{\otimes r_1} \otimes \dots \otimes \mathbf{V}^{\otimes r_n},$$

and

$$\psi = \bigoplus_{R \in \mathbb{N}_0^{\times n}} \psi_R; \quad \psi_R : \mathbf{V}^{\otimes R} \simeq \mathbf{V}^{\otimes R}.$$

¹Since m is unital, and therefore a surjective linear map, any product m' on \mathbf{A} can be obtained in that way. Indeed, $m' = m_\Xi$ with $\Xi = \omega m'$, being ω some right inverse of m .

²A similar construction can be also found in the work of Davidov [3], on twisting of monoidal structures.

By $\mathbb{N}_0^{\times 0}$ we understand the *singleton* set. For the units we write $\mathbb{I}^{\otimes n} = \bigoplus_{R \in \mathbb{N}_0^{\times n}} \mathbb{I}_R$ and $\mathbb{I}_1 = \mathbb{I}$. The composition of two maps, namely ψ and φ , takes the form

$$\psi \varphi = \bigoplus_{R \in \mathbb{N}_0^{\times n}} [\psi \varphi]_R = \bigoplus_{R \in \mathbb{N}_0^{\times n}} \psi_R \varphi_R.$$

This notation lead us to the identification

$$(2.4) \quad \mathbb{C}^n[\mathbf{V}] = \times_{R \in \mathbb{N}_0^{\times n}} \text{Aut}_{\text{Vct}}[\mathbf{V}^{\otimes R}].$$

In other terms, each element of $\mathbb{C}^n[\mathbf{V}]$ can be seen as a section of a fiber bundle with base $\mathbb{N}_0^{\times n}$ and fibers $\text{Aut}_{\text{Vct}}[\mathbf{V}^{\otimes R}]$.

From now on, whenever we are considering a fixed generic vector space \mathbf{V} , we shall write $\mathbb{C}^n[\mathbf{V}] = \mathbb{C}^n$, just for brevity. Based on groups \mathbb{C}^n , we now define a cosimplicial object in the category Grp as follows.

The multiplication m_{\otimes} of \mathbf{V}^{\otimes} restricted to each subspace $\mathbf{V}^{\otimes r} \otimes \mathbf{V}^{\otimes s} \subset \mathbf{V}^{\otimes} \otimes \mathbf{V}^{\otimes}$ is an injective map which has as image the vector subspace $\mathbf{V}^{\otimes r+s} \subset \mathbf{V}^{\otimes}$. Then, m_{\otimes} defines canonical isomorphisms $\mathbf{V}^{\otimes r} \otimes \mathbf{V}^{\otimes s} \simeq \mathbf{V}^{\otimes r+s}$. On the other hand, consider the functions

$$D_i^n = I \times \dots \times D \times \dots \times I : \mathbb{N}_0^{\times n+1} \rightarrow \mathbb{N}_0^{\times n}; \quad i \in \{1, \dots, n\},$$

where the sum on integers $D : (n, m) \mapsto n + m$ acts on the i -th and $i + 1$ -th factors of $\mathbb{N}_0^{\times n+1}$. In terms of these functions, given $R = (r_1, \dots, r_{n+1})$ with $n > 0$, the restrictions to $\mathbf{V}^{\otimes R}$ of the maps

$$(2.5) \quad m_i^n = \underbrace{I \otimes \dots \otimes m_{\otimes}}_{i \text{ factors}} \otimes \underbrace{I \otimes \dots \otimes I}_{n-i \text{ factors}} : (\mathbf{V}^{\otimes})^{\otimes n+1} \rightarrow (\mathbf{V}^{\otimes})^{\otimes n},$$

$i \in \{1, \dots, n\}$, define canonical bijections $m_i^R : \mathbf{V}^{\otimes R} \simeq \mathbf{V}^{\otimes D_i^n(R)}$. The latter give rise to group homomorphisms $\delta_i^n : \mathbb{C}^n \rightarrow \mathbb{C}^{n+1}$ by the assignment

$$(2.6) \quad \psi \mapsto \delta_i^n \psi = \bigoplus_{R \in \mathbb{N}_0^{\times n+1}} [\delta_i^n \psi]_R; \quad [\delta_i^n \psi]_R = (m_i^R)^{-1} \psi_{D_i^n(R)} m_i^R.$$

The elements $\delta_i^n \psi$ are uniquely determined by

$$(2.7) \quad m_i^n \delta_i^n \psi = \psi m_i^n,$$

since the restriction of above equation to $\mathbf{V}^{\otimes R}$ gives precisely the last part of Eq. (2.6). Using the identification $\mathbf{V}^{\otimes R} \approx \mathbf{V}^{\otimes D_i^n(R)}$, we can write in compact form

$$(2.8) \quad \delta_i^n \psi \approx \bigoplus_{R \in \mathbb{N}_0^{\times n+1}} \psi_{D_i^n(R)}.$$

Let us also define the homomorphisms $\delta_{0,n+1}^n : \mathbb{C}^n \rightarrow \mathbb{C}^{n+1}$, $n \neq 0$, as $\delta_0^n \psi = \mathbb{I} \otimes \psi$ and $\delta_{n+1}^n \psi = \psi \otimes \mathbb{I}$. For $n = 0$ any element $\chi \in \mathbb{C}^0$ is a map $1 \mapsto \lambda \in \mathbb{k}^{\times} \equiv \mathbb{k} - \{0\}$. In other words $\mathbb{C}^0 \simeq \mathbb{k}^{\times}$, the multiplicative group of the field \mathbb{k} . So we can define $\delta_{0,1}^0 \chi$ by the assignment $v \in \mathbf{V}^{\otimes} \mapsto \lambda \cdot v \in \mathbf{V}^{\otimes}$. In resume, we have built up a set of group homomorphisms δ_i^n , $i \in \{0, 1, \dots, n+1\}$, $n \in \mathbb{N}_0$. We shall call them *coface operators*.

Consider now, for $i \in \{0, 1, \dots, n\}$, the linear transformations

$$(2.9) \quad \eta_i^n = \underbrace{I \otimes \dots \otimes I}_i \otimes \underbrace{\eta_{\otimes} \otimes \dots \otimes I}_{n+1-i \text{ factors}} : (\mathbf{V}^{\otimes})^{\otimes n} \rightarrow (\mathbf{V}^{\otimes})^{\otimes n+1}$$

related to the unit map $\eta_{\otimes} : \mathbb{k} \rightarrow \mathbf{V}^{\otimes}$, and the functions

$$S_i^n : \mathbb{N}_0^{\times n} \rightarrow \mathbb{N}_0^{\times n+1} \quad / \quad S_i^n(r_1, \dots, r_i, r_{i+1}, \dots, r_n) = (r_1, \dots, r_i, 0, r_{i+1}, \dots, r_n).$$

Since η_{\otimes} as a map over $\mathbf{V}^{\otimes 0} = \mathbb{k}$ is the identity, given $R \in \mathbb{N}_0^{\times n}$ the map η_i^n restricted to $\mathbf{V}^{\otimes R}$ defines a canonical bijection $\eta_i^R : \mathbf{V}^{\otimes R} \simeq \mathbf{V}^{\otimes S_i^n(R)}$. From η_i^n and S_i^n , the group homomorphisms $\sigma_i^n : \mathbb{C}^{n+1} \rightarrow \mathbb{C}^n$, with

$$(2.10) \quad \psi \mapsto \sigma_i^n \psi = \bigoplus_{R \in \mathbb{N}_0^{\times n}} [\sigma_i^n \psi]_R; \quad [\sigma_i^n \psi]_R = (\eta_i^R)^{-1} \psi_{S_i^n(R)} \eta_i^R,$$

can be defined. The equation $\eta_i^n \sigma_i^n \psi = \psi \eta_i^n$ determinates $\sigma_i^n \psi$ completely. From the identification $\mathbf{V}^{\otimes R} \approx \mathbf{V}^{\otimes S_i^n(R)}$ we can write

$$\sigma_i^n \psi \approx \bigoplus_{R \in \mathbb{N}_0^{\times n}} \psi_{S_i^n(R)}.$$

They will be called *codegeneracies*.

Theorem 2. *The map $[n+1] \mapsto \mathbb{C}^n[\mathbf{V}]$ defines a **cosimplicial object** in Grp ; i.e. it can be extended to a functor $\Delta \rightarrow \text{Grp}$, being Δ the simplicial category.*

Proof. We must show that

$$\begin{aligned}
 \delta_i^{n+1} \delta_j^n &= \delta_{j+1}^{n+1} \delta_i^n, & i \leq j; \\
 \sigma_j^{n-1} \sigma_i^n &= \sigma_i^{n-1} \sigma_{j+1}^n, & i \leq j; \\
 \sigma_j^{n+1} \delta_i^{n+1} &= \delta_i^n \sigma_{j-1}^n, & i < j; \\
 \sigma_j^n \delta_i^n &= id, & i = j, i = j + 1; \\
 \sigma_j^{n+1} \delta_i^{n+1} &= \delta_{i-1}^n \sigma_j^n, & i > j + 1.
 \end{aligned}
 \tag{2.11}$$

In what follows, we will omit the index n on maps δ_i^n , σ_i^n , etc. It is easy to see that the opposite equations hold for $\delta = D$, $\sigma = S$ and for $\delta = m$, $\sigma = \eta$. For instance, for $i \leq j$

$$m_j m_i = m_i m_{j+1} \quad \text{and} \quad D_j D_i = D_i D_{j+1}.$$

These are direct consequences of the associative monoidal structures defined by (D, S) in \mathbb{N}_0 and by (m, η) in \mathbf{V}^{\otimes} . Using this fact, let us show the first one of the above equalities. Taking $\psi \in \mathbf{C}^n$ we have for all $R \in \mathbb{N}_0^{\times n+2}$

$$\begin{aligned}
 [\delta_i \delta_j \psi]_R &= [\delta_i (\delta_j \psi)]_R = (m_i^R)^{-1} [\delta_j \psi]_{D_i^{n+1}(R)} m_i^R \\
 &= (m_i^R)^{-1} \left(m_j^{D_i^{n+1}(R)} \right)^{-1} \psi_{D_j^n D_i^{n+1}(R)} m_j^{D_i^{n+1}(R)} m_i^R \\
 &= \left(m_j^{D_i^{n+1}(R)} m_i^R \right)^{-1} \psi_{D_j^n D_i^{n+1}(R)} m_j^{D_i^{n+1}(R)} m_i^R.
 \end{aligned}$$

Note that, for instance, $m_j^{D_i^{n+1}(R)}$ is m_j^{n+1} restricted to $\mathbf{V}^{\otimes D_i^{n+1}(R)}$. Restricting Eq. (2.12) to $\mathbf{V}^{\otimes R}$ and taking into account the domains where m 's are applied, we arrive at

$$\begin{aligned}
 [\delta_i \delta_j \psi]_R &= \left(m_i^{D_{j+1}^{n+1}(R)} m_{j+1}^R \right)^{-1} \psi_{D_i^n D_{j+1}^{n+1}(R)} m_i^{D_{j+1}^{n+1}(R)} m_{j+1}^R \\
 &= (m_{j+1}^R)^{-1} [\delta_i \psi]_{D_{j+1}^{n+1}(R)} m_{j+1}^R = [\delta_{j+1} (\delta_i \psi)]_R = [\delta_{j+1} \delta_i \psi]_R.
 \end{aligned}$$

The others equalities can be shown in an analogous way. ■

Now, we are going to endow the family of groups $\{\mathbf{C}^n[\mathbf{V}]\}_{n \in \mathbb{N}_0}$ with a multiplicative quasicomplex structure, in a very close fashion to the cochain quasicomplex for twisting of bialgebras.

2.2. The multiplicative quasicomplex \mathbf{C}^\bullet . Unless a confusion may arise, the index n on maps δ_i^n will be omitted. Consider the map ∂ that assigns to each $\psi \in \mathbf{C}^n$, for every $n \in \mathbb{N}_0$, an element of \mathbf{C}^{n+1} given by the equation

$$\partial \psi = \left(\prod_{i \text{ odd}}^{\rightarrow} \delta_i \psi \right) \left(\prod_{i \text{ even}}^{\leftarrow} \delta_i \psi \right)^{-1},$$

where $i \in \{0, \dots, n+1\}$ and being

$$\prod_{\{m_1 < \dots < m_k\}}^{\rightarrow} \varphi_i \doteq \varphi_{m_1} \varphi_{m_2} \dots \varphi_{m_k}, \quad \prod_{\{m_1 < \dots < m_k\}}^{\leftarrow} \varphi_i \doteq \varphi_{m_k} \varphi_{m_{k-1}} \dots \varphi_{m_1}.$$

The notation $\partial \psi = \partial_- \psi (\partial_+ \psi)^{-1}$,

$$\partial_- \psi = \prod_{i \text{ odd}}^{\rightarrow} \delta_i \psi, \quad \partial_+ \psi = \prod_{i \text{ even}}^{\leftarrow} \delta_i \psi,$$

will be very useful. Note that $\partial : \mathbf{C}^n \rightarrow \mathbf{C}^{n+1}$ is not a group homomorphism for every n , but satisfies $\partial \mathbb{I}^{\otimes n} = \mathbb{I}^{\otimes n+1}$. The same holds for maps ∂_{\pm} .

Definition 1. We define the **multiplicative cosimplicial quasicomplex of \mathbf{V}** (or its tensor algebra), denoted by the pair $(\mathbf{C}^\bullet[\mathbf{V}], \partial)$, as the sequence

$$\mathbf{C}^0[\mathbf{V}] \xrightarrow{\partial} \mathbf{C}^1[\mathbf{V}] \xrightarrow{\partial} \mathbf{C}^2[\mathbf{V}] \xrightarrow{\partial} \mathbf{C}^3[\mathbf{V}] \xrightarrow{\partial} \dots \quad \blacksquare$$

The prefix *quasi* is used because in general $\partial^2(C^n[\mathbf{V}]) \neq \{\mathbb{I}^{\otimes n+2}\}$. However, we shall see for $n = 0, 1$ the equality holds. Sometimes we shall omit for brevity this prefix.

Let Grp_* be the full subcategory of Set_* whose objects are groups based on their respective unit elements (i.e. its morphisms are unit preserving functions), and denote by Grp_{*q} the category of quasicomplexes in $\text{Grp}_* \subset \text{Set}_*$. Its arrows $f^\bullet : (C^\bullet, \partial_C) \rightarrow (D^\bullet, \partial_D)$ are collections of unit preserving functions $f^n : C^n \rightarrow D^n$ such that equality $f^{n+1} \partial_C = \partial_D f^n$ holds for all n . Then, the pair $(C^\bullet[\mathbf{V}], \partial)$ is an object of Grp_{*q} , and the triple $(C^\bullet[\mathbf{V}], \partial_+, \partial_-)$ is a multiplicative *parity* quasicomplex in Grp_* (see [9] and references therein).

The notions of cocycles, coboundaries, cohomology relation and counitality for (C^\bullet, ∂) are the following.

- An **n-cocycle** is a cochain $\chi \in C^n$ satisfying $\partial\chi = \mathbb{I}^{\otimes n+1}$. We indicate $Z^n = Z^n[\mathbf{V}]$ the set of n -cocycles in C^n .
- Two n -cochains χ and χ' are called **cohomologous** if there exists $\theta \in C^{n-1}$, $n \in \mathbb{N}$, such that the equation $\partial_- \theta \chi (\partial_+ \theta)^{-1} = \chi'$ holds. We denote $\chi \sim_\theta \chi'$ if χ and χ' are cohomologous through θ , or simply $\chi \sim \chi'$, and call $\text{Coh}^n = \text{Coh}^n[\mathbf{V}]$ the subset of $C^n \times C^n$ of cohomologous pairs.
- An **n-coboundary** is an element $\omega \in C^n$ with $\mathbb{I}^{\otimes n} \sim \omega$, i.e. such that $\exists \theta \in C^{n-1}$ satisfying $\partial_- \theta (\partial_+ \theta)^{-1} = \omega$. We denote $B^n = B^n[\mathbf{V}]$ the n -coboundaries of C^n . In particular, $\{\mathbb{I}^{\otimes n}\} \times B^n \subset \text{Coh}^n$, for all $n \in \mathbb{N}$.
- An **n-counital** cochain is a map $\chi \in C^n$ satisfying for $0 \leq i \leq n-1$, $\chi \eta_i^{n-1} = \eta_i^{n-1}$, i.e. $\sigma_i^{n-1} \chi = \mathbb{I}^{\otimes n-1}$. Each $\mathfrak{C}^n = \mathfrak{C}^n[\mathbf{V}]$ indicates the subgroup of counital cochains in C^n .

We must mention that the cohomology relation is not an equivalence relation for all n . In general it is only reflexive, because $\chi \sim \chi$ through $\theta = \mathbb{I}^{\otimes n}$.

As we have said at the beginning of this chapter, we are interested in cochains satisfying Eq. (2.1), i.e. $\chi(\eta \otimes \mathbb{I}) = (\eta \otimes \mathbb{I})$ and $\chi(\mathbb{I} \otimes \eta) = (\mathbb{I} \otimes \eta)$. Since $\eta_0^1 = (\mathbb{I} \otimes \eta)$ and $\eta_1^1 = (\eta \otimes \mathbb{I})$, such conditions can be rewritten $\chi \eta_i^1 = \eta_i^1$, $i = 0, 1$. These are exactly the counital cochains. It follows from (2.11) that, if $\chi \in \mathfrak{C}^n$, then $\delta_i^n \chi$ is not necessarily in \mathfrak{C}^{n+1} . For example, provided $\sigma_i^n \delta_i^n = id$, then $\sigma_i^n \delta_i^n \chi = \chi$. Hence, the groups \mathfrak{C}^n do not define a cosimplicial subobject of C^\bullet . Nevertheless, for the quasicomplex structure we have the following result.

Proposition 1. *The restriction $\partial_{\mathfrak{C}}$ of ∂ to the groups \mathfrak{C}^n defines a quasicomplex $(\mathfrak{C}^\bullet, \partial_{\mathfrak{C}}) \in \text{Grp}_{*q}$, such that the inclusions $u : \mathfrak{C}^n \hookrightarrow C^n$ make \mathfrak{C}^\bullet a subobject of C^\bullet .*

Proof. We must show that $\partial(\mathfrak{C}^n) \subset \mathfrak{C}^{n+1}$. So, let us calculate $\sigma_j^n \partial\psi$ for ψ such that $\sigma_j^{n-1} \psi = \mathbb{I}^{\otimes n-1}$, with $j = 0, \dots, n-1$. Since σ_i^n is a group homomorphism,

$$\sigma_j^n \partial\psi = \left(\prod_{i \text{ odd}}^{\rightarrow} \sigma_j^n \delta_i^n \psi \right) \left(\prod_{i \text{ even}}^{\leftarrow} \sigma_j^n \delta_i^n \psi \right)^{-1}.$$

On the other hand, from (2.11) (remember that $i = 0, 1, \dots, n+1$)

$$\sigma_j^n \delta_i^n \psi = \delta_i^{n-1} \sigma_{j-1}^{n-1} \psi = \delta_i^{n-1} \mathbb{I}^{\otimes n-1} = \mathbb{I}^{\otimes n}, \quad \text{if } i < j;$$

$$\sigma_j^n \delta_i^n \psi = \delta_{i-1}^{n-1} \sigma_j^{n-1} \psi = \delta_{i-1}^{n-1} \mathbb{I}^{\otimes n-1} = \mathbb{I}^{\otimes n}, \quad \text{if } i > j+1;$$

$$\sigma_j^n \delta_i^n \psi = \psi, \quad \text{if } i = j, i = j+1;$$

hence, if j is odd, $\sigma_j^n \partial\psi = \sigma_j^n \delta_j^n \psi (\sigma_j^n \delta_{j+1}^n \psi)^{-1} = \psi \psi^{-1} = \mathbb{I}^{\otimes n}$. The same can be done for j even. Therefore $\sigma_j^n \partial\psi = \mathbb{I}^{\otimes n}$ for $j = 0, \dots, n-1$; that is to say, $\partial\psi \in \mathfrak{C}^{n+1}$ if $\psi \in \mathfrak{C}^n$. ■

From now on, ∂ will denote the coboundary operator for both \mathfrak{C}^\bullet and C^\bullet . It is worth mentioning $(\mathfrak{C}^\bullet, \partial_+, \partial_-)$ is **not** a parity subquasicomplex of $(C^\bullet, \partial_+, \partial_-)$.

The subsets of cocycles, coboundaries and cohomologous cochains in \mathfrak{C}^\bullet will be indicated by $\mathfrak{Z}^n \subset Z^n$, $\mathfrak{B}^n \subset B^n$ and $\mathfrak{Coh}^n \subset \text{Coh}^n$, respectively. Note that $\mathfrak{Z}^n = Z^n \cap \mathfrak{C}^n$, but for \mathfrak{B}^n and \mathfrak{Coh}^n , in general,

$$(2.15) \quad \mathfrak{B}^n \subset B^n \cap \mathfrak{C}^n \quad \text{and} \quad \mathfrak{Coh}^n \subset \text{Coh}^n \cap \mathfrak{C}^n \times \mathfrak{C}^n.$$

2.2.1. *Elementary calculations in low geometric dimensions.*

n = 0 : Given $\chi : \mathbb{k} \simeq \mathbb{k} : 1 \mapsto \lambda$ (so, $\chi^{-1} : 1 \mapsto \lambda^{-1}$), it follows that $\delta_{0,1}\chi : v \mapsto \lambda \cdot v$, and

$$\partial\chi = \delta_1\chi (\delta_0\chi)^{-1} = \mathbb{I} : \mathbf{V}^{\otimes} \rightarrow \mathbf{V}^{\otimes}.$$

Hence $B^1 = \{\mathbb{I}\}$, and as a consequence $B^1 = \mathfrak{B}^1$. In particular, ∂ restricted to C^0 is the trivial group morphism, and accordingly $Z^0 = \{\chi \in C^0 : \partial\chi = \mathbb{I}\} = C^0 = \mathbb{k}^\times$. In particular, $\partial^2\chi = \mathbb{I}^{\otimes 2}$ for every $\chi \in C^0$. On the other hand, the unique counital 0-cochain is the identity map $I_{\mathbb{k}} = \mathbb{I}_0$. Thus, $\mathfrak{C}^0 = \{\mathbb{I}_0\} \subset C^0$.

$\mathbf{n} = \mathbf{1}$: For any $\alpha : \mathbf{V}^\otimes \simeq \mathbf{V}^\otimes$ in \mathcal{C}^1 ,

$$(2.16) \quad \partial\alpha = \delta_1\alpha (\delta_0\alpha)^{-1} (\delta_2\alpha)^{-1} = (\delta_1\alpha) (\alpha \otimes \alpha)^{-1}.$$

Then, a 1-cocycle is an element $\alpha \in \mathcal{C}^1$ such that $\delta_1\alpha = \alpha \otimes \alpha$, or $\alpha_{r+s} \approx \alpha_r \otimes \alpha_s$. In particular, $\alpha_s \approx \alpha_0 \otimes \alpha_s$, so $\alpha_0 = I_{\mathbb{k}}$. This means that α must be counital, i.e. the inclusion $\mathcal{Z}^1 \subset \mathfrak{Z}^1$ holds. Moreover, from Eq. (2.7) the cocycle condition is equivalent to $\alpha m_\otimes = m_\otimes \delta_1\alpha = m_\otimes (\alpha \otimes \alpha)$, hence α is a 1-cocycle *iff* is an algebra automorphism of \mathbf{V}^\otimes obeying $\alpha(\mathbf{V}) \subset \mathbf{V}$. Then, such algebra automorphisms are in bijection with linear automorphisms $Aut_{\mathbf{V}ct}[\mathbf{V}] = GL(n)$, $n = \dim \mathbf{V}$, i.e. $GL(n) \simeq \mathcal{Z}^1 = \mathfrak{Z}^1$. With respect to the square of ∂ on \mathcal{C}^1 , we have

$$\partial^2\alpha = \delta_1(\partial\alpha) \delta_3(\partial\alpha) \delta_0(\partial\alpha)^{-1} \delta_2(\partial\alpha)^{-1} = \delta_1(\partial\alpha) (\partial\alpha \otimes \mathbb{I}) (\mathbb{I} \otimes \partial\alpha)^{-1} \delta_2(\partial\alpha)^{-1},$$

and using that $\delta_1(\alpha \otimes \alpha) = \delta_1\alpha \otimes \alpha$, $\delta_2(\alpha \otimes \alpha) = \alpha \otimes \delta_1\alpha$, and that the δ_i^n 's are group homomorphisms, it follows from (2.16)

$$(2.17) \quad \partial^2\alpha = \delta_1(\delta_1\alpha) \delta_2(\delta_1\alpha)^{-1}.$$

Further, restricting $\delta_{1,2}(\delta_1\alpha)$ to some $\mathbf{V}^{\otimes r} \otimes \mathbf{V}^{\otimes s} \otimes \mathbf{V}^{\otimes t}$, and using the identifications (2.8),

$$[\delta_1(\delta_1\alpha)]_{r,s,t} \approx [\delta_1\alpha]_{r+s,t} \approx \alpha_{r+s+t}, \quad [\delta_2(\delta_1\alpha)]_{r,s,t} \approx [\delta_1\alpha]_{r,s+t} \approx \alpha_{r+s+t},$$

and $[\partial^2\alpha]_{r,s,t} \approx \alpha_{r+s+t} \alpha_{r+s+t}^{-1} = \mathbb{I}_{r,s,t}$ follows, i.e. $\partial^2\alpha = \mathbb{I}^{\otimes 3}$. That means, together with the $n = 0$ case,

$$(2.18) \quad \text{Im } \partial^2|_{\mathcal{C}^{0,1}} = \mathbb{I}^{\otimes 2,3}, \quad \text{or } \mathbf{B}^{1,2} \subset \mathcal{Z}^{1,2}.$$

But ∂ is not a morphism of groups for $n = 1$.

Given $\alpha, \beta \in \mathcal{C}^1$, they are cohomologous *iff* there exists $\lambda \in \mathcal{C}^0 = \mathbb{k}^\times$ such that $\lambda \cdot \alpha \cdot \lambda^{-1} = \beta$ *iff* $\alpha = \beta$. In particular $\mathbf{B}^1 = \{\mathbb{I}\} = \mathfrak{B}^1$, as we have previously seen. Restricting ourself to counital cochains we have $(\alpha, \beta) \in \mathfrak{Coh}^1$ *iff* there exists $\lambda \in \mathcal{C}^0$ such that $\lambda \cdot \alpha \cdot \lambda^{-1} = \beta$. But $\mathcal{C}^0 = \{\mathbb{I}_0\}$, so $(\alpha, \beta) \in \mathfrak{Coh}^1$ *iff* $\alpha = \beta$, and consequently $\mathfrak{Coh}^1 = \text{Coh}^1 \cap \mathcal{C}^1 \times \mathcal{C}^1$.

$\mathbf{n} = \mathbf{2}$: A 2-cocycle is an element $\psi \in \mathcal{C}^2$ satisfying $\delta_1\psi (\psi \otimes \mathbb{I}) = \delta_2\psi (\mathbb{I} \otimes \psi)$, or

$$(2.19) \quad \psi_{r+s,t} (\psi_{r,s} \otimes \mathbb{I}_t) \approx \psi_{r,s+t} (\mathbb{I}_r \otimes \psi_{s,t}).$$

A counital 2-cochain fulfills the equations $\psi(1 \otimes a) = 1 \otimes a$ and $\psi(a \otimes 1) = a \otimes 1$, that is to say, $\psi_{0,s} \approx \mathbb{I}_s$ and $\psi_{r,0} \approx \mathbb{I}_r$. From (2.19) is easy to see that

$$(2.20) \quad \psi_{0,s} \approx \psi_{0,0} \cdot \mathbb{I}_s \quad \text{and} \quad \psi_{r,0} \approx \psi_{0,0} \cdot \mathbb{I}_r,$$

regarding $\psi_{0,0}$ as an element of \mathbb{k}^\times . That means,

$$(2.21) \quad \psi \in \mathfrak{Z}^2 \quad \text{iff} \quad \psi \in \mathcal{Z}^2 \quad \text{and} \quad \psi_{0,0} = 1.$$

Suppose $(\psi, \varphi) \in \text{Coh}^2$, i.e. $\partial_- \theta \psi = \varphi \partial_+ \theta$ for some $\theta \in \mathcal{C}^1$, or in other terms, $\theta_{r+s} \psi_{r,s} \approx \varphi_{r,s} (\theta_r \otimes \theta_s)$. If $(\psi, \varphi) \in \text{Coh}^2 \cap \mathcal{C}^2 \times \mathcal{C}^2$, then $\psi_{0,0} = \varphi_{0,0} = 1$ and $\theta_0 \approx \theta_0 \psi_{0,0} \approx \varphi_{0,0} (\theta_0 \otimes \theta_0) \approx (\theta_0 \otimes \theta_0)$, and in consequence $\theta_0 = 1$, i.e. $\theta \in \mathcal{C}^1$. Therefore, $(\psi, \varphi) \in \mathfrak{Coh}^2$. Summing up (compare with (2.15)),

$$(2.22) \quad \mathfrak{Coh}^i = \text{Coh}^i \cap \mathcal{C}^i \times \mathcal{C}^i \quad \text{and} \quad \mathfrak{B}^i = \mathbf{B}^i \cap \mathcal{C}^i, \quad \text{for } i = 1, 2.$$

2.2.2. (Anti)bicharacter like cochains. Using the terminology of linear forms over a bialgebra [11], we shall say $\psi \in \mathcal{C}^2$ is a *bicharacter* if

$$(2.23) \quad \delta_1\psi = \psi_{23} \psi_{13}, \quad \delta_2\psi = \psi_{12} \psi_{13},$$

and *anti-bicharacter* (see [2] for examples in the bialgebra case) if

$$(2.24) \quad \delta_1\psi = \psi_{13} \psi_{23}, \quad \delta_2\psi = \psi_{13} \psi_{12},$$

where $\psi_{12} = \psi \otimes \mathbb{I}$, $\psi_{23} = \mathbb{I} \otimes \psi$ and $\psi_{13} = (\mathbb{I} \otimes f_\otimes^{-1}) \psi_{12} (\mathbb{I} \otimes f_\otimes)$. We are denoting by f_\otimes the flipping operator on $\mathbf{V}^\otimes \otimes \mathbf{V}^\otimes$. Using the identifications (2.8) and following obvious notation, Eq. (2.23) can be written

$$(2.25) \quad \psi_{r+s,t} \approx (\mathbb{I}_r \otimes \psi_{s,t}) (\mathbb{I}_r \otimes f_{s,t}^{-1}) (\psi_{r,t} \otimes \mathbb{I}_s) (\mathbb{I}_r \otimes f_{s,t}),$$

$$\psi_{r,s+t} \approx (\psi_{r,s} \otimes \mathbb{I}_t) (\mathbb{I}_r \otimes f_{s,t}^{-1}) (\psi_{r,t} \otimes \mathbb{I}_s) (\mathbb{I}_r \otimes f_{s,t}).$$

Note that every (anti)bicharacter ψ is always counital; in fact,

$$\psi_{0,t} = \psi_{0+0,t} \approx (\mathbb{I}_0 \otimes \psi_{0,t}) (\mathbb{I}_0 \otimes f_{0,t}^{-1}) (\psi_{0,t} \otimes \mathbb{I}_0) (\mathbb{I}_0 \otimes f_{0,t}) \approx \psi_{0,t}^2,$$

thus $\psi_{0,t} \approx \mathbb{I}_t$. Also, ψ is completely define by $\psi_{1,1}$.

If ψ is a bicharacter, it will be a 2-cocycle *iff*

$$(2.26) \quad \psi_{12} \psi_{13} \psi_{23} = \psi_{23} \psi_{13} \psi_{12},$$

i.e. ψ satisfies the *Yang-Baxter* (YB) equation. For an anti-bicharacter, ψ is a 2-cocycle iff

$$(2.27) \quad \psi_{12} \psi_{23} = \psi_{23} \psi_{12}.$$

It can be shown it is enough for $\psi_{1,1}$ to satisfy some of the above equations (depending on the case) in order to ensure ψ is a 2-cocycle. Then, given $M \in GL(n^2)$ satisfying (2.26) (resp. (2.27)), we can define a counital 2-cocycle ψ related to an n -dimensional vector space \mathbf{V} , just taking $\psi_{1,1}(a_i \otimes a_j) = M_{ij}^{kl} a_k \otimes a_l$ and extending ψ to all of $\mathbf{V}^{\otimes} \otimes \mathbf{V}^{\otimes}$ using Eq. (2.23) (resp. (2.24)).

2.2.3. First and second cohomology spaces. We have just seen that the cohomology relation in $\mathbf{C}^1[\mathbf{V}]$ is the identity relation, so is an equivalence one. In particular, it can be used to define the *first cohomology space* H^1 as the quotient of \mathbf{Z}^1 , i.e. $H^1 \doteq \mathbf{Z}^1 = \mathfrak{Z}^1 \simeq GL[\mathbf{V}]$. The same can be done for $n = 2$.

Proposition 2. *The cohomology relation for $n = 2$ is an equivalence relation, and every equivalence class with some element in \mathbf{Z}^2 is contained there. This is also valid for \mathfrak{C}^\bullet .*

Proof. We need to show that the relation is symmetric and transitive. Suppose $(\varphi, \psi) \in \text{Coh}^2$, i.e. there exists $\alpha \in \mathbf{C}^1$ such that $\partial_- \alpha \varphi (\partial_+ \alpha)^{-1} = \psi$. Let us first note that ∂_\pm are group homomorphisms when restricted to \mathbf{C}^1 . Indeed, $\partial_- (\alpha) = \delta_1 (\alpha)$ and $\partial_+ (\alpha) = \alpha \otimes \alpha$ for all $\alpha \in \mathbf{C}^1$. In particular $\partial_- (\alpha^{-1}) = \delta_1 (\alpha^{-1}) = (\delta_1 \alpha)^{-1} = (\partial_- \alpha)^{-1}$ and $\partial_+ (\alpha^{-1}) = \alpha^{-1} \otimes \alpha^{-1} = (\partial_+ \alpha)^{-1}$. Thus

$$\partial_- (\alpha^{-1}) \psi \partial_+ (\alpha^{-1})^{-1} = \varphi, \quad \text{i.e. } (\psi, \varphi) \in \text{Coh}^2,$$

and accordingly the relation is symmetric. To show the transitivity, consider again the cohomologous maps φ and ψ , and a 2-cochain $\phi \in \mathbf{C}^2$ such that there exists β satisfying $\partial_- \beta \psi (\partial_+ \beta)^{-1} = \phi$, i.e. $\psi \smile \phi$. Then,

$$\partial_- (\beta \alpha) \varphi (\partial_+ (\alpha \beta))^{-1} = \partial_- \beta \left(\partial_- \alpha \varphi \partial_+ (\alpha)^{-1} \right) (\partial_+ \beta)^{-1} = \phi.$$

Accordingly, $\varphi \smile \phi$ through $\beta \alpha$. It rests to prove each equivalence class containing an element of \mathbf{Z}^2 is a subset of \mathbf{Z}^2 , that is to say, given a 2-cocycle $\varphi \in \mathbf{C}^2$ and a 1-cochain $\alpha \in \mathbf{C}^1$, $\psi = \delta_1 \alpha \varphi (\alpha \otimes \alpha)^{-1}$ is a 2-cocycle too. But,

$$\begin{aligned} \partial_- \psi &= \delta_1 (\delta_1 \alpha) \delta_1 \varphi \left(\delta_1 (\alpha)^{-1} \otimes \alpha^{-1} \right) \left(\delta_1 \alpha \varphi (\alpha \otimes \alpha)^{-1} \otimes \mathbb{I} \right) \\ &= \delta_1 (\delta_1 \alpha) \delta_1 \varphi (\varphi \otimes \mathbb{I}) (\alpha \otimes \alpha \otimes \alpha)^{-1} \end{aligned}$$

and

$$\begin{aligned} (\partial_+ \psi)^{-1} &= \left(\mathbb{I} \otimes (\alpha \otimes \alpha) \varphi^{-1} \delta_1 (\alpha)^{-1} \right) (\alpha \otimes \delta_1 \alpha) \delta_2 (\varphi)^{-1} \delta_2 (\delta_1 \alpha)^{-1} \\ &= (\alpha \otimes \alpha \otimes \alpha) (\mathbb{I} \otimes \varphi^{-1}) \delta_2 (\varphi)^{-1} \delta_2 (\delta_1 \alpha)^{-1}. \end{aligned}$$

Hence, from the last equations and Eq. (2.17)

$$\partial \psi = \delta_1 (\delta_1 \alpha) \partial \varphi \delta_2 (\delta_1 \alpha)^{-1} = \delta_1 (\delta_1 \alpha) \delta_2 (\delta_1 \alpha)^{-1} = \partial^2 \alpha = \mathbb{I}^{\otimes 3},$$

as we wanted to show.

For \mathfrak{C}^\bullet we just have to use the monic $u : \mathfrak{C}^\bullet \hookrightarrow \mathbf{C}^\bullet$, which is given by group monomorphisms. ■

Then we can define the second cohomology space $H^2[\mathbf{V}]$ as the set of cohomologous classes in \mathbf{Z}^2 , i.e. $H^2 = \mathbf{Z}^2 / \sim_{\text{Coh}}$. We are going to see in the next chapter that any 2-cocycle is a 2-coboundary. Therefore, every $\psi \in \mathbf{Z}^2$ is cohomologous to $\mathbb{I}^{\otimes 2}$, and accordingly $H^2[\mathbf{V}] = \{\mathbb{I}^{\otimes 2}\}$. Nevertheless, this does not mean all deformations on a given quantum space generated by \mathbf{V} are trivial, because the maps that define twisted algebras isomorphic to the original one are those cohomologous to the identity through certain class of 1-cochains (what we shall call *admissible*), and not through any of them.

2.3. Functorial properties of $\mathbf{C}^\bullet[\mathbf{V}]$. Let us consider the cosimplicial objects $\mathbf{C}_\mathbf{V}^\bullet : \Delta \rightarrow \text{Grp}$ and $\mathbf{C}_\mathbf{W}^\bullet : \Delta \rightarrow \text{Grp}$ with \mathbf{V} and \mathbf{W} in Vect .

Theorem 3. *The functors $\mathbf{C}_\mathbf{V}^\bullet$ and $\mathbf{C}_\mathbf{W}^\bullet$ are naturally equivalent if \mathbf{V} and \mathbf{W} are isomorphic vector spaces.*

Proof. We must show that there exists a family $\{\alpha_{[n+1]}\}_{n \in \mathbb{N}_0}$ of group isomorphisms

$$\mathbf{C}_\mathbf{V}^\bullet[n+1] = \mathbf{C}^n[\mathbf{V}] \simeq \mathbf{C}^n[\mathbf{W}] = \mathbf{C}_\mathbf{W}^\bullet[n+1]$$

such that

$$\alpha_{[n+2]} \delta_{i,\mathbf{V}}^n = \delta_{i,\mathbf{W}}^n \alpha_{[n+1]} \quad \text{and} \quad \alpha_{[n+1]} \sigma_{i,\mathbf{V}}^n = \sigma_{i,\mathbf{W}}^n \alpha_{[n+2]}.$$

Let f be an isomorphism between \mathbf{V} and \mathbf{W} . Then, the map $f^\otimes : \mathbf{V}^\otimes \rightarrow \mathbf{W}^\otimes$, the unique extension of f to \mathbf{V}^\otimes as an algebra homomorphism, defines for every $n \in \mathbb{N}_0$ a group isomorphism $C^n[\mathbf{V}] \simeq C^n[\mathbf{W}]$ given by

$$(2.28) \quad \psi \mapsto \psi^f \doteq (f^\otimes)^{\otimes n} \psi \left((f^\otimes)^{\otimes n} \right)^{-1}.$$

In particular, since

$$(f^\otimes)^{\otimes n} (\mathbf{V}^{\otimes r_1} \otimes \dots \otimes \mathbf{V}^{\otimes r_n}) \subset \mathbf{W}^{\otimes r_1} \otimes \dots \otimes \mathbf{W}^{\otimes r_n},$$

or in compact form $(f^\otimes)^{\otimes n} (\mathbf{V}^{\otimes R}) \subset \mathbf{W}^{\otimes R}$, we can write

$$(f^\otimes)^{\otimes n} = \bigoplus_{R \in \mathbb{N}_0^{\times n}} f^{\otimes R} \quad \text{and} \quad \psi^f = \bigoplus_{R \in \mathbb{N}_0^{\times n}} f^{\otimes R} \psi_R (f^{\otimes R})^{-1}.$$

Let us call f^n such group isomorphisms, and show that

$$(2.29) \quad f^{n+1} \delta_{i,\mathbf{V}}^n = \delta_{i,\mathbf{W}}^n f^n \quad \text{and} \quad f^{n+1} \sigma_{i,\mathbf{V}}^n = \sigma_{i,\mathbf{W}}^n f^n.$$

Since f^\otimes is an algebra isomorphism, equations

$$(f^\otimes)^{\otimes n} m_{i,\mathbf{V}}^n = m_{i,\mathbf{W}}^n (f^\otimes)^{\otimes n+1} \quad \text{and} \quad (f^\otimes)^{\otimes n+1} \eta_{i,\mathbf{V}}^n = \eta_{i,\mathbf{W}}^n,$$

and analogous equations to its inverse, hold. Restricting to some $R \in \mathbb{N}_0^{\times n+1}$ they translate into

$$(2.30) \quad f^{\otimes D_i^n(R)} m_{i,\mathbf{V}}^R = m_{i,\mathbf{W}}^R f^{\otimes R} \quad \text{and} \quad f^{\otimes S_i^{n+1}(R)} \eta_{i,\mathbf{V}}^R = \eta_{i,\mathbf{W}}^R.$$

For the first equality of (2.29), given $\psi \in C^n[\mathbf{V}]$ and $R \in \mathbb{N}_0^{\times n+1}$, we have

$$[f^{n+1} \delta_{i,\mathbf{V}}^n \psi]_R = f^{\otimes R} [\delta_{i,\mathbf{V}}^n \psi]_R (f^{\otimes R})^{-1} = f^{\otimes R} \left((m_{i,\mathbf{V}}^R)^{-1} \psi_{D_i^n(R)} m_{i,\mathbf{V}}^R \right) (f^{\otimes R})^{-1},$$

and using (2.30),

$$\begin{aligned} [f^{n+1} \delta_{i,\mathbf{V}}^n \psi]_R &= (m_{i,\mathbf{W}}^R)^{-1} f^{\otimes D_i^n(R)} \psi_{D_i^n(R)} (f^{\otimes D_i^n(R)})^{-1} m_{i,\mathbf{W}}^R \\ &= (m_{i,\mathbf{W}}^R)^{-1} \psi_{D_i^n(R)}^f m_{i,\mathbf{W}}^R = [\delta_{i,\mathbf{W}}^n \psi^f]_R = [\delta_{i,\mathbf{W}}^n f^n \psi]_R, \end{aligned}$$

as we wanted to show. The second equality of (2.29) (and the $i = 0, n+1$ cases) can be shown similarly. Thus, taking $\alpha_{[n+1]} = f^n$ we have the natural equivalence $\mathbf{C}_\bullet^{\mathbf{V}} \simeq \mathbf{C}_\bullet^{\mathbf{W}}$ we are looking for. ■

This equivalence can be translated to the corresponding multiplicative quasicomplex structures, as follows. Let us call $\mathcal{G}[\mathbf{Vct}] \subset \mathbf{Vct}$ the groupoid associated to \mathbf{Vct} , i.e. the subcategory of \mathbf{Vct} whose morphisms are isomorphisms.

Theorem 4. *The map $\mathbf{V} \mapsto (\mathbf{C}^\bullet[\mathbf{V}], \partial_{\mathbf{V}})$ defines a functor $\mathcal{G}[\mathbf{Vct}] \rightarrow \mathbf{Grp}_{*q}$.*

Proof. Let us show the map $f \mapsto f^\bullet$, $f : \mathbf{V} \simeq \mathbf{W}$ and $f^n \psi = \psi^f$ for all ψ in $C^n[\mathbf{V}]$ (see Eq. (2.28)), extends $\mathbf{V} \mapsto (\mathbf{C}^\bullet[\mathbf{V}], \partial_{\mathbf{V}})$ to a functor. That is to say, f^\bullet is such that $f^{n+1} \partial_{\mathbf{V}} = \partial_{\mathbf{W}} f^n$. Using Eq. (2.29)

$$(2.31) \quad f^{n+1} \partial_{\mathbf{V}} \psi = \left(\prod_{i \text{ odd}}^{\overleftarrow{}} (f^{n+1} \delta_{i,\mathbf{V}} \psi) \right) \left(\prod_{i \text{ even}}^{\overleftarrow{}} (f^{n+1} \delta_{i,\mathbf{V}} \psi) \right)^{-1}$$

$$= \left(\prod_{i \text{ odd}}^{\overleftarrow{}} \delta_{i,\mathbf{W}} (f^n \psi) \right) \left(\prod_{i \text{ even}}^{\overleftarrow{}} \delta_{i,\mathbf{W}} (f^n \psi) \right)^{-1} = \partial_{\mathbf{W}} f^n \psi,$$

since each f^n is a group homomorphism. Hence, the claim of the proposition follows. ■

Note the functor $\mathcal{G}[\mathbf{Vct}] \rightarrow \mathbf{Grp}_{*q}$ defines an homomorphism of groupoids

$$\mathcal{G}[\mathbf{Vct}] \rightarrow \mathcal{G}[\mathbf{Grp}_{*q}]$$

that assigns to each linear isomorphism $f : \mathbf{V} \simeq \mathbf{W}$ a quasicomplex isomorphism (see (2.28))

$$f^\bullet : (\mathbf{C}^\bullet[\mathbf{V}], \partial_{\mathbf{V}}) \simeq (\mathbf{C}^\bullet[\mathbf{W}], \partial_{\mathbf{W}}) : \psi \mapsto \psi^f.$$

From this, we have the bijections

$$Z^n[\mathbf{V}] \simeq Z^n[\mathbf{W}], \quad B^n[\mathbf{V}] \simeq B^n[\mathbf{W}] \quad \text{and} \quad \text{Coh}^n[\mathbf{V}] \simeq \text{Coh}^n[\mathbf{W}]$$

for all $n \in \mathbb{N}_0$. Clearly, all these results are also true for the subquasicomplexes $\mathbf{e}^\bullet[\mathbf{V}]$ of counital cochains.

In the following subsections we present some structural results. They relate the quasicomplex of a vector space \mathbf{W} with one of its subspaces $\mathbf{V} \subset \mathbf{W}$, and to its dual \mathbf{W}^* . Also relate $\mathbf{C}^\bullet[\mathbf{U}]$ and $\mathbf{C}^\bullet[\mathbf{V}]$ with the

quasicomplexes $\mathbf{C}^\bullet[\mathbf{U} \otimes \mathbf{V}]$. From these results we are able to define a full subcategory of Grp_{*q} , with monoidal structure given by the direct product \times of groups.

All constructions and results will be based on \mathbf{C}^\bullet , but are also valid, at the quasicomplex level, for the related subquasicomplex \mathfrak{C}^\bullet .

2.3.1. Linear subspaces and subobjects in Grp_{*q} . Consider the inclusion of vector spaces $\mathbf{V} \subset \mathbf{W}$. The cochains $\psi \in \mathbf{C}^n[\mathbf{W}]$ such that $\psi(\mathbf{V}^{\otimes R}) \subset \mathbf{V}^{\otimes R}$ holds $\forall R \in \mathbb{N}_0^{\times n}$, form a subgroup of $\mathbf{C}^n[\mathbf{W}]$ that we shall denote $\mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$. Given $\psi \in \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$, let us indicate by $\psi|_{\mathbf{V}}$ and $[\psi|_{\mathbf{V}}]_R$ the restrictions of ψ and ψ_R to $(\mathbf{V}^{\otimes})^{\otimes n}$ and $\mathbf{V}^{\otimes R}$, respectively. Of course, $\psi|_{\mathbf{V}}$ is an element of $\mathbf{C}^n[\mathbf{V}]$ and any element there can be obtained in that way. We are interested on the relation between the cohomological properties of the elements $\psi \in \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$, and the ones of their associated elements $\psi|_{\mathbf{V}}$ in $\mathbf{C}^n[\mathbf{V}]$. To this end, consider the following result.

Proposition 3. *The subgroups $\mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$ define a multiplicative cosimplicial quasicomplex $\mathbf{C}^\bullet[\mathbf{V} \subset \mathbf{W}]$, such that the canonical inclusions and projections*

$$\mathbf{C}^n[\mathbf{V} \subset \mathbf{W}] \hookrightarrow \mathbf{C}^n[\mathbf{W}]; \quad \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}] \rightarrow \mathbf{C}^n[\mathbf{V}] : \psi \mapsto \psi|_{\mathbf{V}},$$

give rise to natural transformations of the respective cosimplicial objects, and to monic and epi arrows, respectively, in Grp_{*q} .

Proof. Note that if $\mathbf{V} \subset \mathbf{W}$ as vector spaces, then $\mathbf{V}^{\otimes} \subset \mathbf{W}^{\otimes}$ as algebras. Hence (see Eq. (2.6)), for every $\psi \in \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$ and $R \in \mathbb{N}_0^{\times n+1}$ we have

$$\begin{aligned} (\delta_{i,\mathbf{W}}\psi)|_{\mathbf{V}^{\otimes R}} &= (m_{i,\mathbf{W}}^R)^{-1} \psi_{D_i^n(R)} m_{i,\mathbf{W}}^R \Big|_{\mathbf{V}^{\otimes R}} \\ (2.32) \quad &= (m_{i,\mathbf{V}}^R)^{-1} [\psi|_{\mathbf{V}}]_{D_i^n(R)} m_{i,\mathbf{V}}^R = [\delta_{i,\mathbf{V}}(\psi|_{\mathbf{V}})]_R. \end{aligned}$$

In particular,

$$\delta_{i,\mathbf{W}}\psi(\mathbf{V}^{\otimes R}) \subset \mathbf{V}^{\otimes R}, \quad \forall R \in \mathbb{N}_0^{\times n+1},$$

i.e. $\delta_{i,\mathbf{W}}\psi \in \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$ if $\psi \in \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$, and consequently Eq. (2.32) can be written

$$(2.33) \quad (\delta_{i,\mathbf{W}}\psi)|_{\mathbf{V}} = \delta_{i,\mathbf{V}}(\psi|_{\mathbf{V}}).$$

Furthermore, the restriction of $\delta_{i,\mathbf{W}}$ to $\mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$ gives a map

$$\delta_{i,\mathbf{V} \subset \mathbf{W}} \doteq \delta_{i,\mathbf{W}}|_{\mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]} : \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}] \rightarrow \mathbf{C}^{n+1}[\mathbf{V} \subset \mathbf{W}],$$

such that

$$(2.34) \quad \delta_{i,\mathbf{V} \subset \mathbf{W}}\psi = \delta_{i,\mathbf{W}}\psi$$

and (using Eq. (2.33))

$$(2.35) \quad (\delta_{i,\mathbf{V} \subset \mathbf{W}}\psi)|_{\mathbf{V}} = \delta_{i,\mathbf{V}}(\psi|_{\mathbf{V}}).$$

The same is true for the maps $\sigma_{i,\mathbf{W}}$. In consequence, from $\delta_{i,\mathbf{V} \subset \mathbf{W}}$ and $\sigma_{i,\mathbf{V} \subset \mathbf{W}}$ a cosimplicial object

$$\mathbf{C}_{\mathbf{V} \subset \mathbf{W}}^\bullet : \mathbf{\Delta} \rightarrow \text{Grp} : [n+1] \mapsto \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$$

is defined and, since Eqs. (2.34) and (2.35), the proposition follows. ■

We must mention the group injections $\mathbf{C}^n[\mathbf{V}] \hookrightarrow \mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$ given by the map $\phi \mapsto \phi \oplus \mathbb{I}_{\mathbf{K}}$, where we are decomposing the algebra $(\mathbf{W}^{\otimes})^{\otimes n}$ in the subalgebra $(\mathbf{V}^{\otimes})^{\otimes n}$ and a bilateral ideal \mathbf{K} , are not morphisms of quasicomplexes (the problem is with the coface operators $\delta_{0,n+1}^n$). Hence, the resulting group monomorphisms $\mathbf{C}^n[\mathbf{V}] \hookrightarrow \mathbf{C}^n[\mathbf{W}]$ do not define $\mathbf{C}^\bullet[\mathbf{V}]$ as a subobject of $\mathbf{C}^\bullet[\mathbf{W}]$.

Now, to the wanted result. The cohomological properties of the elements $\psi \in \mathbf{C}^\bullet[\mathbf{V} \subset \mathbf{W}]$, determinate completely the ones of the restrictions $\psi|_{\mathbf{V}} \in \mathbf{C}^\bullet[\mathbf{V}]$. More precisely,

Proposition 4. *Let ψ, φ be elements of $\mathbf{C}^n[\mathbf{V} \subset \mathbf{W}]$.*

a) $\psi \in \mathbf{Z}^n[\mathbf{V} \subset \mathbf{W}]$ iff $\psi \in \mathbf{Z}^n[\mathbf{W}]$ iff $\psi|_{\mathbf{V}} \in \mathbf{Z}^n[\mathbf{V}]$.

b) If $(\psi, \varphi) \in \text{Coh}^n[\mathbf{V} \subset \mathbf{W}]$, then $(\psi, \varphi) \in \text{Coh}^n[\mathbf{W}]$ and $(\psi|_{\mathbf{V}}, \varphi|_{\mathbf{V}}) \in \text{Coh}^n[\mathbf{V}]$.

Proof. a) The first part follows from the facts that $\partial_{\mathbf{V} \subset \mathbf{W}}\psi = \partial_{\mathbf{W}}\psi$, and

$$(\partial_{\mathbf{V} \subset \mathbf{W}}\psi)|_{\mathbf{V}} = \partial_{\mathbf{W}}\psi|_{\mathbf{V}} = \partial_{\mathbf{V}}(\psi|_{\mathbf{V}})$$

(direct consequences of Eqs. (2.34) and (2.35), resp.).

b) $(\psi, \varphi) \in \text{Coh}^n[\mathbf{V} \subset \mathbf{W}]$ means that there exists $\theta \in \mathbf{C}^{n-1}[\mathbf{V} \subset \mathbf{W}]$ such that

$$[\partial_{\mathbf{V} \subset \mathbf{W}}]_- \theta \psi = \varphi [\partial_{\mathbf{V} \subset \mathbf{W}}]_+ \theta.$$

It is clear this implies $(\psi, \varphi) \in \text{Coh}^n[\mathbf{W}]$. In addition, $\theta|_{\mathbf{V}}$, $[\partial_{\mathbf{V} \subset \mathbf{W}}]_- \theta|_{\mathbf{V}}$ and $[\partial_{\mathbf{V} \subset \mathbf{W}}]_+ \theta|_{\mathbf{V}}$ define elements of $\mathbf{C}^\bullet[\mathbf{V}]$, thus

$$\partial_{\mathbf{V},-}(\theta|_{\mathbf{V}})(\psi|_{\mathbf{V}}) = ([\partial_{\mathbf{V} \subset \mathbf{W}}]_- \theta)|_{\mathbf{V}} \psi|_{\mathbf{V}} = [\partial_{\mathbf{V} \subset \mathbf{W}}]_- \theta \psi|_{\mathbf{V}} = \varphi [\partial_{\mathbf{V} \subset \mathbf{W}}]_+ \theta|_{\mathbf{V}} = (\varphi|_{\mathbf{V}}) \partial_{\mathbf{V},+}(\theta|_{\mathbf{V}}),$$

i.e. $\psi|_{\mathbf{V}} \smile \varphi|_{\mathbf{V}}$ through $\theta|_{\mathbf{V}}$. ■

2.3.2. Coadjoints and products. Let us restrict ourself to the category $\text{Vct}_f \subset \text{Vct}$ of finite dimensional \mathbb{k} -vector spaces. In such a case, for each isomorphism $\mathbf{V} \simeq \mathbf{V}^*$ there is a related isomorphism of quasicomplexes $(\mathbf{C}^\bullet[\mathbf{V}], \partial_{\mathbf{V}}) \simeq (\mathbf{C}^\bullet[\mathbf{V}^*], \partial_{\mathbf{V}^*})$ and, in particular, group isomorphisms $\mathbf{C}^n[\mathbf{V}] \simeq \mathbf{C}^n[\mathbf{V}^*]$ for every n . On the other hand, the natural pairing between $(\mathbf{V}^\otimes)^{\otimes n}$ and $(\mathbf{V}^{*\otimes})^{\otimes n}$, namely

$$\langle \cdot, \cdot \rangle : (\mathbf{V}^{*\otimes})^{\otimes n} \times (\mathbf{V}^\otimes)^{\otimes n} \rightarrow \mathbb{k},$$

gives rise to group anti-homomorphisms (the *transposition maps*)

$$*_l : \mathbf{C}^n[\mathbf{V}] \rightarrow \mathbf{C}^n[\mathbf{V}^*] : \psi \mapsto \psi^*$$

with $\langle \psi^* v, w \rangle = \langle v, \psi w \rangle$, and

$$*_r : \mathbf{C}^n[\mathbf{V}^*] \rightarrow \mathbf{C}^n[\mathbf{V}] : \psi \mapsto \psi^*,$$

being $\langle v, \psi^* w \rangle = \langle \psi v, w \rangle$. In both cases we have $(\psi^*)_R = (\psi_R)^*$, $\forall R$, and consequently,

$$(2.36) \quad \delta_{\mathbf{V}^*,i} \psi^* = (\delta_{i,\mathbf{V}} \psi)^* \quad \text{and} \quad \sigma_{\mathbf{V}^*,i} \psi^* = (\sigma_{i,\mathbf{V}} \psi)^*.$$

Of course, $*_l *_r = *_r *_l = id$; that is to say, $*_l$ and $*_r$ define mutually inverse natural equivalences of the corresponding cosimplicial objects. But these maps do not define morphisms between quasicomplexes $(\mathbf{C}^\bullet[\mathbf{V}], \partial_{\mathbf{V}})$ and $(\mathbf{C}^\bullet[\mathbf{V}^*], \partial_{\mathbf{V}^*})$, unless they are composed with the inversion $\psi \mapsto \psi^{-1}$.

Proposition 5. *The maps $\psi \in \mathbf{C}^n[\mathbf{V}] \mapsto \psi^{*-1} = \psi^{-1*}$, $n \in \mathbb{N}_0$, define an isomorphism in Grp_{*q} . The same is true for $\psi \in \mathbf{C}^n[\mathbf{V}^*] \mapsto \psi^{*-1}$.*

Proof. Since they are clearly bijective maps, we just must prove that

$$(\partial_{\mathbf{V}} \psi)^{*-1} = \partial_{\mathbf{V}^*}(\psi^{*-1}), \quad \forall \psi \in \mathbf{C}^n[\mathbf{V}], \quad n \in \mathbb{N}_0.$$

Using Eq. (2.36), we have

$$\begin{aligned} (\partial_{\mathbf{V}} \psi)^* &= \left(\overleftarrow{\prod}_{i \text{ odd}} \delta_{i,\mathbf{V}} \psi \overleftarrow{\prod}_{i \text{ even}} (\delta_{i,\mathbf{V}} \psi)^{-1} \right)^* = \overleftarrow{\prod}_{i \text{ even}} (\delta_{i,\mathbf{V}} \psi)^{*-1} \overleftarrow{\prod}_{i \text{ odd}} (\delta_{i,\mathbf{V}} \psi)^* \\ &= \overleftarrow{\prod}_{i \text{ even}} (\delta_{\mathbf{V}^*,i} \psi^*)^{-1} \overleftarrow{\prod}_{i \text{ odd}} \delta_{\mathbf{V}^*,i} \psi^* = \overleftarrow{\prod}_{i \text{ even}} \delta_{\mathbf{V}^*,i}(\psi^{*-1}) \overleftarrow{\prod}_{i \text{ odd}} \delta_{\mathbf{V}^*,i} \psi^*, \end{aligned}$$

where in the last step we use δ 's are homomorphisms of groups. Inverting the last member we arrive at the wanted result, i.e.

$$(\partial_{\mathbf{V}} \psi)^{*-1} = \overleftarrow{\prod}_{i \text{ odd}} \delta_{\mathbf{V}^*,i}(\psi^{*-1}) \overleftarrow{\prod}_{i \text{ even}} (\delta_{\mathbf{V}^*,i}(\psi^{*-1}))^{-1} = \partial_{\mathbf{V}^*}(\psi^{*-1}).$$

In the same way the another claim can be shown. ■

For reason that will become clear later, we indicate ψ^{*-1} by $\psi^!$. Following this notation, we can call $\mathbf{C}^\bullet[\mathbf{V}^*]$ the *coadjoint* quasicomplex to $\mathbf{C}^\bullet[\mathbf{V}]$, and denote it $\mathbf{C}^\bullet[\mathbf{V}]^!$. Its coboundary is $\partial_{\mathbf{V}^*} \psi = \partial_{\mathbf{V}}(\psi^!)$. Given $g : \mathbf{C}^\bullet[\mathbf{V}] \rightarrow \mathbf{C}^\bullet[\mathbf{W}]$, its coadjoint arrow $g^! : \mathbf{C}^\bullet[\mathbf{V}]^! \rightarrow \mathbf{C}^\bullet[\mathbf{W}]^!$ can be defined as $g^!(\psi) = (g(\psi^!))^!$. Of course, $g^!$ is (by composition) a morphism in Grp_{*q} and, $\mathbf{C}^\bullet[\mathbf{V}]^{!!} = \mathbf{C}^\bullet[\mathbf{V}]$, $g^{!!} = g$.

In these terms, if $\psi \in \mathbf{Z}^n[\mathbf{V}]$, we have as a consequence of the last proposition that $\psi^! \in \mathbf{Z}^n[\mathbf{V}]^!$. In addition, if ψ is an (anti)bicharacter in $\mathbf{C}^2[\mathbf{V}]$, then $\psi^!$ is an (anti)bicharacter in $\mathbf{C}^2[\mathbf{V}]^!$.

We also can define a product between our quasicomplexes. Indeed, for every $n \in \mathbb{N}_0$, and $\mathbf{V}, \mathbf{W} \in \text{Vct}$, consider the groups $\mathbf{C}^n[\mathbf{V}] \times \mathbf{C}^n[\mathbf{W}]$. We shall indicate $\mathbf{C}^\bullet[\mathbf{V}] \times \mathbf{C}^\bullet[\mathbf{W}]$ the *product cosimplex* of $\mathbf{C}^\bullet[\mathbf{V}]$ and $\mathbf{C}^\bullet[\mathbf{W}]$, with object function $[n+1] \mapsto \mathbf{C}^n[\mathbf{V}] \times \mathbf{C}^n[\mathbf{W}]$ and coface operators and codegeneracies

$$(2.37) \quad \delta_i^n(\phi, \varphi) = (\delta_{i,\mathbf{V}}^n \phi, \delta_{i,\mathbf{W}}^n \varphi), \quad \sigma_i^n(\phi, \varphi) = (\sigma_{i,\mathbf{V}}^n \phi, \sigma_{i,\mathbf{W}}^n \varphi).$$

Naturally, the associated *product quasicomplex* will have a coboundary map such that $\partial(\phi, \varphi) = (\partial_{\mathbf{V}} \phi, \partial_{\mathbf{W}} \varphi)$. It is clear that given a pair of morphisms $f^\bullet : \mathbf{C}^\bullet[\mathbf{V}] \rightarrow \mathbf{C}^\bullet[\mathbf{V}']$ and $g^\bullet : \mathbf{C}^\bullet[\mathbf{W}] \rightarrow \mathbf{C}^\bullet[\mathbf{W}']$, the related maps

$$f^n \times g^n : \mathbf{C}^n[\mathbf{V}] \times \mathbf{C}^n[\mathbf{W}] \rightarrow \mathbf{C}^n[\mathbf{V}'] \times \mathbf{C}^n[\mathbf{W}']$$

define an arrow $\mathbf{C}^\bullet[\mathbf{V}] \times \mathbf{C}^\bullet[\mathbf{W}] \rightarrow \mathbf{C}^\bullet[\mathbf{V}'] \times \mathbf{C}^\bullet[\mathbf{W}']$.

Proposition 6. $\mathbf{C}^\bullet[\mathbf{V}] \times \mathbf{C}^\bullet[\mathbf{W}]$ is a subobject of $\mathbf{C}^\bullet[\mathbf{V} \otimes \mathbf{W}]$ in Grp_{*q} .

Proof. Consider the canonical isomorphisms $t_R : (\mathbf{V} \otimes \mathbf{W})^{\otimes R} \simeq \mathbf{V}^{\otimes R} \otimes \mathbf{W}^{\otimes R}$. We shall show the group monomorphisms $j : \mathbf{C}^n[\mathbf{V}] \times \mathbf{C}^n[\mathbf{W}] \hookrightarrow \mathbf{C}^n[\mathbf{V} \otimes \mathbf{W}]$,

$$j(\phi, \varphi) = \bigoplus_{R \in \mathbb{N}_0^{x_n}} [j(\phi, \varphi)]_R = \bigoplus_{R \in \mathbb{N}_0^{x_n}} t_R^{-1} (\phi_R \otimes \varphi_R) t_R,$$

define a morphism in Grp_{*q} . To see that, it is enough to show $j \delta_i^n(\phi, \varphi) = \delta_{i, \mathbf{V} \otimes \mathbf{W}}^n j(\phi, \varphi)$. Given $R \in \mathbb{N}_0^{x_n+1}$, by definition of j ,

$$\begin{aligned} [j \delta_i^n(\phi, \varphi)]_R &= t_R^{-1} \left([\delta_{i, \mathbf{V}}^n \phi]_R \otimes [\delta_{i, \mathbf{W}}^n \varphi]_R \right) t_R \\ &= t_R^{-1} \left((m_{i, \mathbf{V}}^R)^{-1} \phi_{D_i^n(R)} m_{i, \mathbf{V}}^R \otimes (m_{i, \mathbf{W}}^R)^{-1} \varphi_{D_i^n(R)} m_{i, \mathbf{W}}^R \right) t_R \\ &= t_R^{-1} (m_{i, \mathbf{V}}^R \otimes m_{i, \mathbf{W}}^R)^{-1} \left(\phi_{D_i^n(R)} \otimes \varphi_{D_i^n(R)} \right) (m_{i, \mathbf{V}}^R \otimes m_{i, \mathbf{W}}^R) t_R. \end{aligned}$$

Noting that $m_{i, \mathbf{V} \otimes \mathbf{W}}^R = t_{D_i^n(R)}^{-1} (m_{i, \mathbf{V}}^R \otimes m_{i, \mathbf{W}}^R) t_R$, the last member is precisely

$$(m_{i, \mathbf{V} \otimes \mathbf{W}}^R)^{-1} [j(\phi, \varphi)]_{D_i^n(R)} m_{i, \mathbf{V} \otimes \mathbf{W}}^R = [\delta_{i, \mathbf{V} \otimes \mathbf{W}}^n j(\phi, \varphi)]_R.$$

The same can be done for $\delta_{0, n+1}^n$. This concludes our proof. ■

In forthcoming sections, we shall identify $(\mathbf{V} \otimes \mathbf{W})^{\otimes R}$ with $\mathbf{V}^{\otimes R} \otimes \mathbf{W}^{\otimes R}$, enabling us to describe each $j(\phi, \varphi)$ as a map

$$(2.38) \quad j(\phi, \varphi) = \bigoplus_{R \in \mathbb{N}_0^{x_n}} [j(\phi, \varphi)]_R = \bigoplus_{R \in \mathbb{N}_0^{x_n}} \phi_R \otimes \varphi_R,$$

or equivalently, as the restriction of $\phi \otimes \varphi$ to

$$\bigoplus_{R \in \mathbb{N}_0^{x_n}} \mathbf{V}^{\otimes R} \otimes \mathbf{W}^{\otimes R} \subset (\mathbf{V}^{\otimes})^{\otimes n} \otimes (\mathbf{W}^{\otimes})^{\otimes n},$$

i.e. $j(\phi, \varphi) \subset \phi \otimes \varphi$.

Let us denote $\mathbf{C}^\bullet[\text{Vct}]$ and $\mathbf{C}^\bullet[\text{Vct}_f]$ the full subcategories of Grp_{*q} whose objects are of the form

$$\mathbf{C}^\bullet\{\mathbf{V}_i\} \doteq \mathbf{C}^\bullet[\mathbf{V}_1] \times \dots \times \mathbf{C}^\bullet[\mathbf{V}_m]; \quad \mathbf{V}_i \in \text{Vct},$$

and define $\mathbf{C}^\bullet\{\mathbf{V}_i\}^! \doteq \mathbf{C}^\bullet[\mathbf{V}_1]^! \times \dots \times \mathbf{C}^\bullet[\mathbf{V}_m]^!$ for $\mathbf{V}_i \in \text{Vct}_f$. The following result resumes all we have discussed.

Theorem 5. Each quasicomplex $\mathbf{C}^\bullet\{\mathbf{V}_i\}$ is a subobject of $\mathbf{C}^\bullet[\otimes_i \mathbf{V}_i]$. The map

$$\times : (\mathbf{C}^\bullet\{\mathbf{V}_i\}, \mathbf{C}^\bullet\{\mathbf{W}_i\}) \mapsto \mathbf{C}^\bullet\{\mathbf{V}_i\} \times \mathbf{C}^\bullet\{\mathbf{W}_i\}$$

defines a monoidal structure in $\mathbf{C}^\bullet[\text{Vct}]$ with unit $\mathbf{C}^\bullet[\mathbb{k}]$; while $! : \mathbf{C}^\bullet\{\mathbf{V}_i\} \mapsto \mathbf{C}^\bullet\{\mathbf{V}_i\}^!$ gives rise to a covariant monoidal functor in $\mathbf{C}^\bullet[\text{Vct}_f]$. ■

It is important to note, among other things, the set inclusions (under natural identifications)

$$(2.39) \quad Z^n[\mathbf{V}_1] \times \dots \times Z^n[\mathbf{V}_m] \subset Z^n[\mathbf{V}_1 \otimes \dots \otimes \mathbf{V}_m]$$

hold. This is a consequence of the following more general fact. Since the image of $\mathcal{G}[\text{Vct}] \rightarrow \text{Grp}_{*q}$ (see §2.3), is contained in $\mathbf{C}^\bullet[\text{Vct}]$, it can be regarded as a functor $\mathbf{C}^\bullet : \mathcal{G}[\text{Vct}] \rightarrow \mathbf{C}^\bullet[\text{Vct}]$. Then, the monics $j : \mathbf{C}^\bullet[\mathbf{V}] \times \mathbf{C}^\bullet[\mathbf{W}] \hookrightarrow \mathbf{C}^\bullet[\mathbf{V} \otimes \mathbf{W}]$ define natural transformations $\mathbf{C}^\bullet \times \mathbf{C}^\bullet \rightarrow \mathbf{C}^\bullet(\cdot \otimes \cdot)$. The same is true for the full subfunctor $\mathbf{C}_f^\bullet : \mathcal{G}[\text{Vct}_f] \rightarrow \mathbf{C}^\bullet[\text{Vct}_f]$. On the other hand, for each $\mathbf{V} \in \text{Vct}_f$, $\mathbf{C}^\bullet[\mathbf{V}] \simeq \mathbf{C}^\bullet[\mathbf{V}^*]$ is a functorial isomorphism for the natural equivalences $\mathbf{C}_f^\bullet \simeq \mathbf{C}_f^\bullet *$ and $\mathbf{C}_f^\bullet \simeq ! \mathbf{C}_f^\bullet$.

2.4. Comparison with the quicomplex for bialgebra twisting. Let \mathbf{A} be a bialgebra, i.e. $\mathbf{A} \in \mathbb{k}\text{-Bialg} = \text{Bialg}$, with coalgebra structure (Δ, ε) . Consider the groups $\mathbf{G}^n[\mathbf{A}]$, $n \in \mathbb{N}_0$, formed out by the invertible elements of $\text{Lin}[\mathbf{A}^{\otimes n}, \mathbb{k}]$ under the convolution product $*$. Such a product is given by

$$\psi * \varphi = (\psi \otimes \varphi) \Delta^{(n)}, \quad \psi, \varphi \in \text{Lin}[\mathbf{A}^{\otimes n}, \mathbb{k}],$$

being $\Delta^{(n)}$ the usual coproduct on $\mathbf{A}^{\otimes n}$ (in particular, $\Delta^{(1)} = \Delta$ and $\Delta^{(0)} = I_{\mathbb{k}}$). The unit of $*$ is $\varepsilon^{\otimes n}$, the usual counit of $\mathbf{A}^{\otimes n}$.

Denote by the pair (m, η) the algebra structure of \mathbf{A} , and define (as we have done for a tensor algebra \mathbf{V}^{\otimes}) the maps m_i^n and η_i^n as in Eqs. (2.5) and (2.9). Then, the group homomorphisms $d_i^n : \mathbf{G}^n[\mathbf{A}] \rightarrow \mathbf{G}^{n+1}[\mathbf{A}]$ and $s_i^n : \mathbf{G}^{n+1}[\mathbf{A}] \rightarrow \mathbf{G}^n[\mathbf{A}]$, with

$$d_i^n : \psi \mapsto \begin{cases} \varepsilon \otimes \psi, & \text{if } i = 0, \\ \psi m_i^n, & \text{if } i \in \{1, \dots, n\}, \\ \psi \otimes \varepsilon, & \text{if } i = n + 1, \end{cases}$$

and

$$s_i^n : \psi \mapsto \psi \eta_i^n, \quad i \in \{0, 1, \dots, n\},$$

define a cosimplicial object $\mathbf{G}_{\mathbf{A}}^{\bullet} : [n+1] \mapsto \mathbf{G}^n[\mathbf{A}]$ in $\text{Grp}[4][11]$. Moreover, using the convolution product, the map

$$(2.40) \quad d : \psi \mapsto \left(\prod_{i \text{ even}}^{\rightarrow} d_i \psi \right)^{-1} * \left(\prod_{i \text{ odd}}^{\leftarrow} d_i \psi \right),$$

where $i \in \{0, \dots, n+1\}$, supplies the groups $\mathbf{G}^n[\mathbf{A}]$ with a (cosimplicial multiplicative) cochain quicomplex structure.³ Note that the order is reversed w.r.t. the coboundary ∂ of the quicomplexes \mathbf{C}^{\bullet} .

We recall that $\psi \in \mathbf{G}^n[\mathbf{A}]$ is a counital n -cochain if satisfies $s_i^n \psi = \psi \eta_i^n = \varepsilon^{\otimes n}$ for all i . Also recall, a bicharacter is a map $\psi \in \mathbf{G}^2[\mathbf{A}]$ such that

$$d_1 \psi = \psi(m \otimes I) = \psi_{13} * \psi_{23}, \quad d_2 \psi = \psi(I \otimes m) = \psi_{13} * \psi_{12},$$

while an anti-bicharacter [2] fulfill the opposite equations: $d_1 \psi = \psi_{23} * \psi_{13}$ and $d_2 \psi = \psi_{12} * \psi_{13}$.

Since d is a unit preserving map, the pair $(\mathbf{G}^{\bullet}[\mathbf{A}], d)$ is a quicomplex in Grp_* , and the factorization of d in d_+ , d_- , namely $d\psi = (d_+ \psi)^{-1} d_- \psi$, makes the triple $(\mathbf{G}^{\bullet}[\mathbf{A}], d_+, d_-)$ a parity quicomplex there. For later convenience, we mention the following known facts without proof.

Proposition 7. *If a pair of bialgebras \mathbf{A} and \mathbf{B} are homomorphic by means of a map $\alpha : \mathbf{A} \rightarrow \mathbf{B}$, then there exists a natural transformation $\mathbf{G}_{\mathbf{B}}^{\bullet} \rightarrow \mathbf{G}_{\mathbf{A}}^{\bullet}$ of cosimplicial objects given by functions*

$$(2.41) \quad \psi \in \mathbf{G}^n[\mathbf{B}] \mapsto \psi \alpha^{\otimes n} \in \mathbf{G}^n[\mathbf{A}].$$

Moreover, the same functions define a morphism $(\mathbf{G}^{\bullet}[\mathbf{B}], d_{\mathbf{B}}) \rightarrow (\mathbf{G}^{\bullet}[\mathbf{A}], d_{\mathbf{A}})$ of quicomplexes, becoming the assignment $\mathbf{A} \mapsto \mathbf{G}^{\bullet}[\mathbf{A}]$ into a contravariant functor $\text{Bialg} \rightarrow \text{Grp}_{*q}$. ■

When the bialgebra \mathbf{A} is a tensor algebra \mathbf{C}^{\otimes} ,⁴ we can ask which is the relation between $\mathbf{G}^{\bullet}[\mathbf{C}^{\otimes}]$ and $\mathbf{C}^{\bullet}[\mathbf{C}]$. In order to find this relation, given $\psi \in \mathbf{G}^n[\mathbf{C}^{\otimes}]$, consider the endomorphism of $(\mathbf{C}^{\otimes})^{\otimes n}$ of the form⁵

$$\psi * \mathbb{I}^{\otimes n} * \psi^{-1} = (\psi \otimes \mathbb{I}^{\otimes n} \otimes \psi^{-1}) \left(\mathbb{I}^{\otimes n} \otimes \Delta^{(n)} \right) \Delta^{(n)}.$$

Given another $\varphi \in \mathbf{G}^n[\mathbf{C}^{\otimes}]$, we have

$$(\psi * \varphi) * \mathbb{I}^{\otimes n} * (\psi * \varphi)^{-1} = \psi * \varphi * \mathbb{I}^{\otimes n} * \varphi^{-1} * \psi^{-1} = (\varphi * \mathbb{I}^{\otimes n} * \varphi^{-1}) (\psi * \mathbb{I}^{\otimes n} * \psi^{-1}),$$

and obviously for $\varepsilon^{\otimes n}$, $\varepsilon^{\otimes n} * \mathbb{I}^{\otimes n} * \varepsilon^{\otimes n} = \mathbb{I}^{\otimes n}$. It follows that $\psi^{-1} * \mathbb{I}^{\otimes n} * \psi$ is the inverse of $\psi * \mathbb{I}^{\otimes n} * \psi^{-1}$, hence the latter is a linear automorphism of $(\mathbf{C}^{\otimes})^{\otimes n}$. Therefore, the function $F^n : \psi \mapsto \psi^{-1} * \mathbb{I}^{\otimes n} * \psi$ is a group **anti**-homomorphism, i.e. $F^n(\varepsilon^{\otimes n}) = \mathbb{I}^{\otimes n}$ and $F^n(\psi * \varphi) = F^n \varphi F^n \psi$. Suppose from now on the comultiplication on \mathbf{C}^{\otimes} is such that the inclusion $\Delta(\mathbf{C}) \subset \mathbf{C} \otimes \mathbf{C}$ holds. In other words, we are saying \mathbf{C} is a coalgebra. Then, since $\Delta^{(n)}$ is an algebra morphism, for every $R \in \mathbb{N}_0^{\times n}$

$$\text{Im } \Delta_R \subset \mathbf{C}^{\otimes R} \otimes \mathbf{C}^{\otimes R}, \quad \Delta_R \doteq \Delta^{(n)} \Big|_{\mathbf{C}^{\otimes R}},$$

³We reorder the factor, with respect to [11], in a way that is convenient for our work.

⁴Every tensor algebra \mathbf{C}^{\otimes} has, fixing a basis $\{v_i\}$ on \mathbf{C} , a bialgebra structure given by the extensions of $\Delta : v_i \mapsto v_i \otimes v_i$ and $\varepsilon : v_i \mapsto 1$ to algebra homomorphisms. This is cocommutative.

⁵Given a bialgebra \mathbf{A} and two linear maps $\alpha, \beta : \mathbf{A} \rightarrow \mathbf{B}_{\alpha, \beta}$, $\alpha * \beta$ will always mean $(\alpha \otimes \beta) \Delta : \mathbf{A} \rightarrow \mathbf{B}_{\alpha} \otimes \mathbf{B}_{\beta}$.

and accordingly each map $F^n\psi$ is homogeneous, thus an n -cochain in $C^n[\mathbf{C}]$. Moreover,

Theorem 6. *Let \mathbf{C} be a coalgebra. Then there exist group anti-homomorphisms $F^n : G^n[\mathbf{C}^\otimes] \rightarrow C^n[\mathbf{C}]$, $n \in \mathbb{N}_0$, that define:*

- a natural transformation of the functors $G_{\mathbf{C}^\otimes}^\bullet$ and $C_{\mathbf{C}}^\bullet$, and
- a morphism of cochain quasicomplexes $(G^\bullet[\mathbf{C}^\otimes], d) \rightarrow (C^\bullet[\mathbf{C}], \partial)$ in Grp_{*q} .

Proof. For the first statement we must show that $F^{n+1}d_i^n = \delta_i^n F^n$ and $F^{n+1}s_i^n = \sigma_i^n F^{n+2}$. Let us apply the first member of the first equality to $\psi \in G^n[\mathbf{C}^\otimes]$. We have, for $i \neq 0, n+1$, the element of $C^{n+1}[\mathbf{C}]$

$$\begin{aligned} F^{n+1}d_i^n\psi &= F^{n+1}(\psi m_i^n) = (\psi m_i^n) * \mathbb{I}^{\otimes n+1} * (\psi^{-1} m_i^n) \\ &= (\psi m_i^n \otimes \mathbb{I}^{\otimes n+1} \otimes \psi^{-1} m_i^n) \left(\mathbb{I}^{\otimes n+1} \otimes \Delta^{(n+1)} \right) \Delta^{(n+1)}, \end{aligned}$$

which restricted to $\mathbf{C}^{\otimes R}$, for some $R \in \mathbb{N}_0^{\times n+1}$, gives

$$[F^{n+1}d_i^n\psi]_R = (\psi m_i^R \otimes \mathbb{I}_R \otimes \psi^{-1} m_i^R) (\mathbb{I}_R \otimes \Delta_R) \Delta_R.$$

Writing $\mathbb{I}_R = (m_i^R)^{-1} m_i^R$ and using

$$(2.42) \quad (m_i^R \otimes m_i^R) \Delta_R = \Delta_{D_i^{n+1}(R)} m_i^R,$$

it follows that $[F^{n+1}d_i^n\psi]_R$ is equal to

$$\begin{aligned} &(m_i^R)^{-1} \left[(\psi \otimes \mathbb{I}_{D_i^{n+1}(R)} \otimes \psi^{-1}) \left(\mathbb{I}_{D_i^{n+1}(R)} \otimes \Delta_{D_i^{n+1}(R)} \right) \Delta_{D_i^{n+1}(R)} \right] m_i^R \\ &= (m_i^R)^{-1} [\psi * \mathbb{I}^{\otimes n} * \psi^{-1}]_{D_i^{n+1}(R)} m_i^R = [\delta_i^n F^n\psi]_R. \end{aligned}$$

If for instance $i = 0$, then⁶

$$\begin{aligned} F^{n+1}(\varepsilon \otimes \psi) &= (\varepsilon \otimes \psi) * \mathbb{I}^{\otimes n+1} * (\varepsilon \otimes \psi^{-1}) = (\varepsilon \otimes \psi) * (\mathbb{I} \otimes \mathbb{I}^{\otimes n}) * (\varepsilon \otimes \psi^{-1}) \\ &= (\varepsilon * \mathbb{I} * \varepsilon) \otimes (\psi * \mathbb{I}^{\otimes n} * \psi^{-1}) = \mathbb{I} \otimes F^n\psi. \end{aligned}$$

The other equalities can be shown analogously. We must use for $F^{n+1}s_i^n = \sigma_i^n F^{n+2}$ that

$$(2.43) \quad \varepsilon_R = \varepsilon_{D_i^{n+1}(R)} m_i^R,$$

what follows from the algebra map character of ε .

Let us prove the second statement. $F^\bullet : G^\bullet[\mathbf{C}^\otimes] \rightarrow C^\bullet[\mathbf{C}]$ is a morphism in Grp_{*q} if and only if equation $F^{n+1}d = \partial F^n$ is satisfied. Using $F^{n+1}d_i^n = \delta_i^n F^n$ and the fact that each F^n is an anti-homomorphism of groups, the last equation is immediately fulfilled (compare Eqs. (2.13) with (2.40)). That concludes our proof. ■

The above theorem describes completely the relation between $G^\bullet[\mathbf{C}^\otimes]$ and $C^\bullet[\mathbf{C}]$ (provided \mathbf{C} is a coalgebra) as multiplicative cosimplicial quasicomplexes. Nevertheless, let us look a little closer at the maps F^n . Observe that

$$F^n\psi = \mathbb{I}^{\otimes n} \text{ iff } \psi * \mathbb{I}^{\otimes n} = \mathbb{I}^{\otimes n} * \psi$$

iff $\psi * \varphi = \varphi * \psi$ for all $\varphi \in G^n[\mathbf{C}^\otimes]$. In other words, $F^n\psi = \mathbb{I}^{\otimes n}$ iff $\psi \in \mathcal{Z}^n$, the center of $G^n[\mathbf{C}^\otimes]$. In particular, if the coproduct Δ of \mathbf{C}^\otimes is cocommutative, then $F^n\psi = \mathbb{I}^{\otimes n}$ for all ψ in $G^n[\mathbf{C}^\otimes]$, for all n . In any case, we may consider the quotient groups $G_{\mathcal{Z}}^n[\mathbf{C}^\otimes] \doteq G^n[\mathbf{C}^\otimes]/\mathcal{Z}^n$ and the corresponding injections⁷ $i^n : G_{\mathcal{Z}}^n[\mathbf{C}^\otimes] \hookrightarrow C^n[\mathbf{C}]$ induced by each F^n . Thus, roughly speaking, the elements of G^n form, up to the center, a subset of C^n . Moreover, we have:

Theorem 7. *The groups $G_{\mathcal{Z}}^n[\mathbf{C}^\otimes]$ define a multiplicative cosimplicial quasicomplex in such a way that the commutative diagrams of group (anti)homomorphisms*

$$\begin{array}{ccc} & G_{\mathcal{Z}}^n[\mathbf{C}^\otimes] & \\ p^n \nearrow & & \searrow i^n \\ G^n[\mathbf{C}^\otimes] & \xrightarrow{F^n} & C^n[\mathbf{C}] \end{array}$$

⁶We are using that given a pair of coalgebras $(\mathbf{A}, \Delta_{\mathbf{A}})$ and $(\mathbf{B}, \Delta_{\mathbf{B}})$, and linear maps α, β and γ, δ with domain \mathbf{A} and \mathbf{B} , resp., the convolution product related to the usual coalgebra in $\mathbf{A} \otimes \mathbf{B}$ satisfies $(\alpha \otimes \gamma) * (\beta \otimes \delta) = (\alpha * \beta) \otimes (\gamma * \delta)$.

⁷As we will see later, from the point of view of twisting of bialgebras, the groups of interest are precisely $G_{\mathcal{Z}}^n$, instead of G^n .

where $p^n : \mathbf{G}^n[\mathbf{C}^\otimes] \rightarrow \mathbf{G}_{\mathcal{Z}}^n[\mathbf{C}^\otimes]$ is the canonical projection, give rise to a commutative diagram of natural transformations among their related cosimplicial objects, and a commutative diagram of morphisms in Grp_{*q} of their corresponding cochain quasicomplex. ■

The theorem follows from the lemmas below. To enunciate the first one, let us note that given $R \in \mathbb{N}_0^{\times n}$, the pair $(\Delta_R, \varepsilon_R)$ defines a coalgebra structure on $\mathbf{C}^{\otimes R}$. Then, the set $\text{Lin}[\mathbf{C}^\otimes]$ is supplied with a convolution product $*$ which enable us to define the groups $\mathbf{G}^R = \mathbf{G}^R[\mathbf{C}^\otimes]$, given by the invertible linear forms $\mathbf{C}^{\otimes R} \rightarrow \mathbb{k}$. Of course, the unit is ε_R . On the other hand, because of Eq. (2.3), every element ψ in $\mathbf{G}^n = \mathbf{G}^n[\mathbf{C}^\otimes]$ is defined by a family of linear forms $\psi_R \in \mathbf{G}^R$, one for each $R \in \mathbb{N}_0^{\times n}$, being $\psi_R = \psi|_{\mathbf{C}^{\otimes R}}$. In other words, the elements of \mathbf{G}^n can be characterized as functions $R \in \mathbb{N}_0^{\times n} \mapsto \psi_R \in \mathbf{G}^R$. Furthermore, the restrictions $\psi \mapsto \psi_R$ are group epimorphisms, since $[\psi * \varphi]_R = \psi_R * \varphi_R$.

Lemma 1. *The following statements are equivalent:*

- i) $\psi \in \mathcal{Z}^n \subset \mathbf{G}^n$.
- ii) The equation $\psi_R * \gamma = \gamma * \psi_R$ holds $\forall \gamma \in \mathbf{G}^R, \forall R \in \mathbb{N}_0^{\times n}$.

Proof. A linear form ψ is in the center of \mathbf{G}^n iff $\psi_R * \varphi_R = \varphi_R * \psi_R$ for all $R \in \mathbb{N}_0^{\times n}$, for all $\varphi \in \mathbf{G}^n$. But, to give an arbitrary element $\varphi \in \mathbf{G}^n$ is the same as giving for each R an arbitrary element of \mathbf{G}^R , by the characterization above. Then, $\psi \in \mathcal{Z}^n$ iff $\psi_R * \gamma = \gamma * \psi_R$ for all $\gamma \in \mathbf{G}^R$ and $R \in \mathbb{N}_0^{\times n}$. ■

Lemma 2. *For all $n \in \mathbb{N}_0$, $d_i^n(\mathcal{Z}^n) \subset \mathcal{Z}^{n+1}$ and $s_i^n(\mathcal{Z}^{n+1}) \subset \mathcal{Z}^n$.*

Proof. We shall only prove the inclusions $d_i^n(\mathcal{Z}^n) \subset \mathcal{Z}^{n+1}$, since the another ones can be proven in a similar way. Let ψ and φ be arbitrary linear forms in \mathbf{G}^n and \mathbf{G}^{n+1} , respectively. Then, for $i \neq 0, n+1$,

$$(d_i^n \psi) * \varphi = (\psi m_i^n \otimes \varphi) \Delta^{(n+1)}$$

restricted to some $\mathbf{C}^{\otimes R}$ ($R \in \mathbb{N}_0^{\times n+1}$) is equal to

$$\begin{aligned} (\psi m_i^R \otimes \varphi) \Delta_R &= \left(\psi \otimes \varphi (m_i^R)^{-1} \right) (m_i^R \otimes m_i^R) \Delta_R \\ &= \left((\psi \otimes \gamma) \Delta_{D_i^{n+1}(R)} \right) m_i^R = \left(\psi_{D_i^{n+1}(R)} * \gamma \right) m_i^R, \end{aligned}$$

where $\gamma \doteq \varphi (m_i^R)^{-1}$ is a linear form $\mathbf{C}^{\otimes D_i^{n+1}(R)} \rightarrow \mathbb{k}$. This linear form is invertible w.r.t. the convolution product, and $\gamma^{-1} \doteq \varphi^{-1} (m_i^R)^{-1}$, because (see Eqs. (2.42) and (2.43))

$$\gamma^{-1} * \gamma = \left(\varphi^{-1} (m_i^R)^{-1} \otimes \varphi (m_i^R)^{-1} \right) \Delta_{D_i^{n+1}(R)} = (\varphi^{-1} \otimes \varphi) \Delta_R (m_i^R)^{-1} = \varepsilon_R (m_i^R)^{-1} = \varepsilon_{D_i^{n+1}(R)}.$$

Therefore, $\gamma \in \mathbf{G}^{D_i^{n+1}(R)}$. Now, if $\psi \in \mathcal{Z}^n$, from the last lemma

$$\left(\psi_{D_i^{n+1}(R)} * \gamma \right) m_i^R = \left(\gamma * \psi_{D_i^{n+1}(R)} \right) m_i^R = (\varphi \otimes \psi m_i^R) \Delta_R,$$

for all R , hence $(d_i^n \psi) * \varphi = \varphi * (d_i^n \psi)$ for all $\varphi \in \mathbf{G}^{n+1}$. In the $i = 0$ case,

$$(\varepsilon \otimes \psi) * \varphi = \varphi [(\varepsilon \otimes \psi) * (\mathbb{I} \otimes \mathbb{I}^{\otimes n})] = \varphi [(\varepsilon * \mathbb{I}) \otimes (\psi * \mathbb{I}^{\otimes n})] = \varphi [\mathbb{I} \otimes (\psi * \mathbb{I}^{\otimes n})],$$

and similarly

$$\varphi * (\varepsilon \otimes \psi) = \varphi [(\mathbb{I} * \varepsilon) \otimes (\mathbb{I}^{\otimes n} * \psi)] = \varphi [\mathbb{I} \otimes (\mathbb{I}^{\otimes n} * \psi)].$$

In addition, we have supposed $\psi \in \mathcal{Z}^n$, then $\psi * \mathbb{I}^{\otimes n} = \mathbb{I}^{\otimes n} * \psi$ and $d_0^n \psi * \varphi = \varphi * d_0^n \psi$ holds. And analogously for $i = n+1$. Hence, $d_i^n(\mathcal{Z}^n) \subset \mathcal{Z}^{n+1}$ for all n, i . ■

As a direct consequence of the last theorem, the cocycles, (anti)bicharacters, coboundaries and every cohomologous class of $\mathbf{G}_{\mathcal{Z}}^\bullet$ can be seen as subsets of the cocycles, the coboundaries and corresponding cohomologous class of \mathbf{C}^\bullet , respectively. Furthermore, the counital cochains of $\mathbf{G}_{\mathcal{Z}}^\bullet$ define another quasicomplex, namely $\mathfrak{G}_{\mathcal{Z}}^\bullet$, such that there exists a monic $\mathfrak{G}_{\mathcal{Z}}^\bullet \hookrightarrow \mathbf{C}^\bullet$ given by group anti-monomorphisms.

By last, consider a vector space \mathbf{V} and the free bialgebra generated by a *multiplicative matrix* $t_i^j = v^j \otimes v_i \in \mathbf{V}^* \otimes \mathbf{V}$. That is to say, the algebra is $[\mathbf{V}^* \otimes \mathbf{V}]^\otimes$ and the coalgebra structure is given by assignments $\Delta : t_i^j \mapsto t_i^k \otimes t_k^j$ and $\varepsilon : t_i^j \mapsto \delta_i^j$. It can be shown the bijection $\text{End}[\mathbf{V}] = \text{Lin}[\mathbf{V}, \mathbf{V}] \simeq \text{Lin}[\mathbf{V}^* \otimes \mathbf{V}, \mathbb{k}]$ gives rise to an isomorphism of quasicomplexes

$$(2.44) \quad \mathbf{C}^\bullet[\mathbf{V}] \simeq \mathbf{G}^\bullet \left[[\mathbf{V}^* \otimes \mathbf{V}]^\otimes \right].$$

3. TWISTING OF QUANTUM LINEAR SPACES

As a previous step toward the quantum space twist transformations we need the concept of *admissible* cochain. Consider a pair $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ and a 2-cochain $\psi \in \mathcal{C}^2[\mathbf{A}_1]$. The map ψ defines a linear map in $\mathbf{A} \otimes \mathbf{A}$ iff

$$(3.1) \quad \psi(\mathbf{A}_1^{\otimes} \otimes \ker \Pi + \ker \Pi \otimes \mathbf{A}_1^{\otimes}) \subset \mathbf{A}_1^{\otimes} \otimes \ker \Pi + \ker \Pi \otimes \mathbf{A}_1^{\otimes},$$

where Π is the canonical epimorphism $\mathbf{A}_1^{\otimes} \rightarrow \mathbf{A}$. Such a map, namely $\Psi : \mathbf{A} \otimes \mathbf{A} \rightarrow \mathbf{A} \otimes \mathbf{A}$, would be given by the equation $\Psi \Pi^{\otimes 2} = \Pi^{\otimes 2} \psi$.

Definition 2. We shall call \mathcal{A} -*admissible*, or *admissible* for \mathcal{A} , the n -cochains satisfying

$$\psi(\ker \Pi^{\otimes n}) \subset \ker \Pi^{\otimes n} \quad \text{and} \quad \psi^{-1}(\ker \Pi^{\otimes n}) \subset \ker \Pi^{\otimes n},$$

or equivalently, $\psi(\ker \Pi^{\otimes n}) = \ker \Pi^{\otimes n}$. ■

If \mathcal{A} is conic, admissibility condition for $\psi \in \mathcal{C}^2[\mathbf{A}_1]$ reduces to

$$(3.2) \quad \psi(\mathbf{A}_1^{\otimes r} \otimes \mathbf{I}_s + \mathbf{I}_r \otimes \mathbf{A}_1^{\otimes s}) \subset \mathbf{A}_1^{\otimes r} \otimes \mathbf{I}_s + \mathbf{I}_r \otimes \mathbf{A}_1^{\otimes s},$$

for all $r, s \in \mathbb{N}_0$, where $\ker \Pi = \bigoplus_{n \geq 2} \mathbf{I}_n$. The other inclusions (and therefore the equalities) follow from the fact that the involved vector spaces are finite dimensional.

The \mathcal{A} -admissible n -cochains form a subgroup of $\mathcal{C}^n[\mathbf{A}_1]$. It can be shown the quotient of this subgroup by the identification

$$(3.3) \quad \psi \sim \psi' \Leftrightarrow \Pi^{\otimes n}(\psi - \psi') = 0,$$

is such that the map $[\psi] \mapsto \Psi$, from admissible equivalence classes to $\text{Aut}_{\text{Vct}}[\mathbf{A}^{\otimes n}]$, is an injective group homomorphism.

It is worth remarking admissibility of ψ does not imply the one of $\partial\psi$. For instance, consider the algebra generated by $\mathbf{A}_1 = \text{span}[a, b]$ and quotient by the relation $ab = 0$. Define the 1-cochain $\theta : \mathbf{A}_1^{\otimes} \simeq \mathbf{A}_1^{\otimes}$ such that θ_n is the identity unless for the elements abx and xab , where $\theta(abx) = xab$ and $\theta(xab) = abx$, being x equal to a or b . Hence, θ is clearly admissible. Then, for instance,

$$\delta_1 \theta(ab \otimes b) = (m_1^R)^{-1} \theta_3 m_1^R(ab \otimes b) = (m_1^R)^{-1}(bab) = ba \otimes b,$$

with $R = (2, 1)$, what implies $\delta_1 \theta(\ker \Pi^{\otimes 2}) \neq \ker \Pi^{\otimes 2}$, i.e. $\partial\theta$ is not admissible. Therefore, unfortunately the subgroups of admissible cochains do not define a subcomplex of $\mathcal{C}^\bullet[\mathbf{A}_1]$. Nevertheless, in §4.1 we consider for each conic quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ a stronger condition than admissibility, which enable us to define a subcomplex $\mathcal{C}^\bullet[\mathcal{A}] \subset \mathcal{C}^\bullet[\mathbf{A}_1]$.

3.1. Counital 2-cocycle twisting. Consider a quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ with related unital associative algebra structure (m, η) , and an admissible counital 2-cochain $\psi \in \mathcal{C}^2[\mathbf{A}_1]$. Since ψ gives rise to a unique automorphism Ψ of $\mathbf{A} \otimes \mathbf{A}$ by the equation $\Psi \Pi^{\otimes 2} = \Pi^{\otimes 2} \psi$, we can define, as at the beginning of §2, a new algebra structure over \mathbf{A} , namely $\mathbf{A}_\psi \doteq (\mathbf{A}, m_\psi)$, with $m_\psi = m \Psi$. In these terms we have the following notion of twisting process on quantum spaces.

Definition 3. Given a quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A}) \in \text{FGA}$ (resp. CA) and a related admissible $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$, we define the *twisting of \mathcal{A} by ψ* as the pair $\mathcal{A}_\psi = (\mathbf{A}_1, \mathbf{A}_\psi) \in \text{FGA}$ (resp. CA). ■

The following theorem justifies above definition.

Theorem 8. Let Ψ be a map defined by an \mathcal{A} -admissible counital 2-cocycle $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$. Then $m_\psi = m \Psi$ is an associative product with unit η , and the associative unital algebra $\mathbf{A}_\psi \doteq (\mathbf{A}, m_\psi, \eta)$ is also generated by \mathbf{A}_1 . Moreover, m_ψ defines a filtration that coincides with the one associated to m (see Eq. (1.1)). In particular, for the conic case, m_ψ defines the same gradation as m .

Proof. Since Π is a morphism of unital algebras $\Pi m_\otimes = m \Pi$ and $\Pi(\lambda) = \eta(\lambda)$, $\forall \lambda \in \mathbf{A}_1^{\otimes 0} = \mathbb{k}$. Then, if ψ is counital we have

$$\Psi(\eta(\lambda) \otimes [a]) = \Psi \Pi^{\otimes 2}(\lambda \otimes a) = \Pi^{\otimes 2} \psi(\lambda \otimes a) = \eta(\lambda) \otimes [a],$$

being $[a] = \Pi(a)$. So, $\Psi(\eta \otimes I) = (\eta \otimes I)$ and

$$m_\psi(\eta \otimes I) = m \Psi(\eta \otimes I) = m(\eta \otimes I) = I.$$

The same holds for $I \otimes \eta$; hence η is a unit map for the product m_ψ . Let us see that m_ψ is associative. It will be associative iff

$$m_\psi(m_\psi \otimes I) \Pi^{\otimes 3} = m_\psi(I \otimes m_\psi) \Pi^{\otimes 3},$$

iff

$$\Pi m_\otimes \psi(m_\otimes \otimes I)(\psi \otimes I) = \Pi m_\otimes \psi(I \otimes m_\otimes)(I \otimes \psi).$$

From Equation (2.7) for $n = 2$, $\psi(m_\otimes \otimes I) = (m_\otimes \otimes I)\delta_1\psi$ and $\psi(I \otimes m_\otimes) = (I \otimes m_\otimes)\delta_2\psi$, so m_ψ is associative iff

$$\Pi m_\otimes (I \otimes m_\otimes) (\partial\psi - \mathbb{I}^{\otimes 3}) = 0,$$

where we have used the associativity of m_\otimes . But the last eq. holds trivially since ψ is a 2-cocycle.

It rests to show that \mathbf{A}_1 generates \mathbf{A} through m_ψ . We know that, given a basis $\{a_i\}$ of \mathbf{A}_1 , it is possible to express any element of \mathbf{A} as a linear combination of words written with the *letters* a_i 's and *glued* with the product m . Let us show the same is true for m_ψ . We will proceed by induction on the *length* of the words.

Denoting m by a blank space and m_ψ by “ \cdot ”, the map Ψ for $r, s \in \mathbb{N}_0$ is given by a $(\dim \mathbf{A}_1)^{r+s}$ square invertible matrix representing $\psi_{r,s}$, namely (remember that $\Pi(a_i) = a_i$)

$$\Psi(a_{i_1} \dots a_{i_r} \otimes a_{j_1} \dots a_{j_s}) = \psi_{i_1 \dots i_r, j_1 \dots j_s}^{k_1 \dots k_r, l_1 \dots l_s} a_{k_1} \dots a_{k_r} \otimes a_{l_1} \dots a_{l_s}.$$

Since

$$a_i \cdot a_j = m_\psi(a_i \otimes a_j) = \psi_{i,j}^{k,l} m(a_k \otimes a_l) = \psi_{i,j}^{k,l} a_k a_l,$$

we have

$$a_k a_l = (\psi^{-1})_{k,l}^{i,j} a_i \cdot a_j.$$

Then, for words of length 2 there exists an invertible linear map $\theta_2 : \mathbf{A}_1^{\otimes 2} \simeq \mathbf{A}_1^{\otimes 2}$, with matrix coefficients $(\theta_2)_{ij}^{kl} = \psi_{i,j}^{k,l}$, such that

$$a_k a_l = (\theta_2^{-1})_{ij}^{kl} a_i \cdot a_j.$$

Suppose that for words of n letters there exists an invertible linear map θ_n such that

$$(3.4) \quad a_{i_1} \dots a_{i_n} = (\theta_n^{-1})_{i_1 \dots i_n}^{j_1 \dots j_n} a_{j_1} \cdot \dots \cdot a_{j_n}.$$

Then, for $n+1$

$$\begin{aligned} (a_{j_1} \cdot \dots \cdot a_{j_n}) \cdot a_{j_{n+1}} &= (\theta_n)_{j_1 \dots j_n}^{i_1 \dots i_n} (a_{i_1} \dots a_{i_n}) \cdot a_{j_{n+1}} \\ &= (\theta_n)_{j_1 \dots j_n}^{i_1 \dots i_n} \psi_{i_1 \dots i_n, j_{n+1}}^{k_1 \dots k_r, k_{n+1}} a_{k_1} \dots a_{k_{n+1}} \doteq (\theta_{n+1})_{j_1 \dots j_{n+1}}^{k_1 \dots k_{n+1}} a_{k_1} \dots a_{k_{n+1}}, \end{aligned}$$

i.e.

$$a_{k_1} \dots a_{k_{n+1}} = (\theta_{n+1}^{-1})_{k_1 \dots k_{n+1}}^{j_1 \dots j_{n+1}} a_{j_1} \cdot \dots \cdot a_{j_{n+1}},$$

being

$$(3.5) \quad (\theta_{n+1}^{-1})_{k_1 \dots k_{n+1}}^{j_1 \dots j_{n+1}} \doteq (\psi^{-1})_{k_1 \dots k_n, k_{n+1}}^{i_1 \dots i_n, j_{n+1}} (\theta_n^{-1})_{i_1 \dots i_n}^{j_1 \dots j_n}.$$

Hence, by induction, any element of \mathbf{A} can be expressed as linear combinations of words constructed with elements of \mathbf{A}_1 and glued with m_ψ . Both words have the same length, in the sense that can be obtained as the image by Π of homogeneous elements with same number of factors. From that it is clear m_ψ defines the same filtration as m , and for the conic case the same gradation, as we wanted to show. \blacksquare

In other words, the last theorem says the pair $(\mathbf{A}_1, \mathbf{A}_\psi)$ is a quantum space, and if $(\mathbf{A}_1, \mathbf{A})$ is conic, so is its twisting. In the following, we shall describe some interesting examples.

3.1.1. *The quantum plane as a 2-cocycle twisting.* Let $\mathcal{A}^{2|0} = (\mathbf{A}_1, \mathbf{A})$ be the quantum space with $\mathbf{A}_1 = \text{span}[a, b]$ and $\mathbf{A} = \mathbb{k}[a, b]$, the commutative algebra freely generated by the symbols a and b . In other words, $\mathcal{A}^{2|0}$ gives the coordinate ring of the two dimensional affine space. Consider the basis

$$\{a^{f_1} b^{g_1} \dots a^{f_k} b^{g_k} : f, g \in \mathbb{N}_0, k \in \mathbb{N}\}$$

of \mathbf{A}_1^{\otimes} , and for every element v in that basis the positive numbers $m^a(v) \doteq \sum_{i=1}^k f_i$ and $m^b(v) \doteq \sum_{i=1}^k g_i$. Now, given $\hbar \in \mathbb{k}$ let us define the 2-cochain

$$\psi_\hbar : v \otimes w \mapsto \exp[m^a(v) m^b(w) \hbar] v \otimes w.$$

The counitality is immediate, since taking $v = w = 1$, equation $m^a(v) m^b(w) = 0$ holds. This cochain is \mathcal{A} -admissible, in fact, ψ_\hbar gives rise to a linear automorphism Ψ_\hbar of $\mathbf{A}^{\otimes 2}$ such that

$$(3.6) \quad \Psi_\hbar(a^n b^m \otimes a^r b^s) = \exp(ns\hbar) a^n b^m \otimes a^r b^s.$$

The formula above defines Ψ_\hbar completely, because the set $\{a^n b^m\}_{n, m \in \mathbb{N}_0}$ is a basis for \mathbf{A} . Let us see that ψ_\hbar is a 2-cocycle. Straightforward calculations show that

$$\delta_1 \psi_\hbar(u \otimes v \otimes w) = \exp[(m^a(u) + m^a(v)) m^b(w) \hbar] u \otimes v \otimes w,$$

$$\delta_2 \psi_\hbar(u \otimes v \otimes w) = \exp[m^a(u) (m^b(v) + m^b(w)) \hbar] u \otimes v \otimes w,$$

for any elements u, v and w of the basis of \mathbf{A}_1^{\otimes} . Now, applying consecutively $\psi_{\hbar} \otimes \mathbb{I}$ and $\delta_1 \psi_{\hbar}$ on such elements we have

$$\begin{aligned} u \otimes v \otimes w &\mapsto \exp [m^a(u) m^b(v) \hbar] u \otimes v \otimes w \\ &\mapsto \exp [(m^a(u) + m^a(v)) m^b(w) \hbar] \exp [m^a(u) m^b(v) \hbar] u \otimes v \otimes w \\ &= \exp [(m^a(u) m^b(w) + m^a(v) m^b(w) + m^a(u) m^b(v)) \hbar] u \otimes v \otimes w, \end{aligned}$$

and doing the same for $\mathbb{I} \otimes \psi_{\hbar}$ and $\delta_2 \psi_{\hbar}$,

$$\begin{aligned} u \otimes v \otimes w &\mapsto \exp [m^a(v) m^b(w) \hbar] u \otimes v \otimes w \\ &\mapsto \exp [m^a(u) (m^b(v) + m^b(w)) \hbar] \exp [m^a(v) m^b(w) \hbar] u \otimes v \otimes w \\ &= \exp [(m^a(u) m^b(v) + m^a(u) m^b(w) + m^a(v) m^b(w)) \hbar] u \otimes v \otimes w. \end{aligned}$$

Thus $\delta_1 \psi_{\hbar} (\psi_{\hbar} \otimes \mathbb{I}) = \delta_2 \psi_{\hbar} (\mathbb{I} \otimes \psi_{\hbar})$. Moreover, it can be seen that ψ_{\hbar} is a bicharacter and also an anti-bicharacter. Accordingly, the product $m_{\psi_{\hbar}} = m \Psi_{\hbar} = \star$ is associative, and from Eq. (3.6) for $n = s = 0$ and $m = r = 1$, $a \star b = \exp \hbar b \star a$ follows. Thus, $\mathcal{A}_{\psi_{\hbar}}^{2|0}$ is isomorphic to the quantum plane

$$\mathcal{A}_{\hbar}^{2|0} \doteq \mathbb{k} \langle x, y \rangle / \{xy - \exp \hbar yx\}$$

under the extension of $x \mapsto a, y \mapsto b$ to an algebra map. The map ψ_{\hbar} also defines an admissible 2-cocycle for the superplane $\mathcal{A}^{0|2}$, given by a Grassmann algebra in two variables. Writing again $m_{\psi_{\hbar}} = \star$, we have the relation $a \star b = -\exp \hbar b \star a$, thus $\mathcal{A}_{\psi_{\hbar}}^{0|2}$ is isomorphic to the quantum 0|2-dimensional superplane

$$\mathcal{A}_{-\hbar}^{0|2} = \mathbb{k} \langle \eta, \xi \rangle / \{\eta \xi + \exp \hbar \xi \eta; \eta^2; \xi^2\}.$$

This result can easily be generalized to any quantum space $\mathcal{A}_{\hbar}^{n|0}$ and $\mathcal{A}_{\hbar}^{0|n}$. In general, given $M \in GL(n^2)$ fulfilling the YB equation or such that $M_{12} M_{23} = M_{23} M_{12}$, we can define, as at the end of §2.2.2, a bicharacter or an anti-bicharacter ψ related to an n -dimensional vector space \mathbf{A}_1 . Such a cochain is always admissible for commutative and anticommutative quantum spaces \mathcal{A} generated by \mathbf{A}_1 . The ideal related to \mathcal{A}_{ψ} is generated by

$$(3.7) \quad (N_{ij}^{kl} \mp N_{ji}^{kl}) a_k \otimes a_l; \quad N = M^{-1},$$

the \mp corresponding to the \mathcal{A} commutative or \mathcal{A} anticommutative cases, respectively. (We need the results of §3.2.2 to show that.) Note relations in (3.7) can be written $(R_{ij}^{kl} \mp P_{ij}^{kl}) a_k \otimes a_l$, being $R = M_{21} N$ and P the permutation matrix. It can be shown R is a YB operator, in fact is a *triangular* one, i.e. $R_{21} R = I$. Thus, twisting \mathcal{A} by (anti)bicharacters define quantum spaces with relations given by triangular YB operators of the form $M_{21} M^{-1}$. As a particular case, to obtain $\mathcal{A}_{\hbar}^{n|0}$ or $\mathcal{A}_{-\hbar}^{0|n}$ from \mathcal{A} , ψ can be defined by

$$M_{ij}^{kl} = \exp \{ [1 - \text{sg}(i - j)] \hbar / 2 \} \delta_i^k \delta_j^l; \quad i, j, k, l = 1 \dots n;$$

being $\text{sg}(0) = 1$. The extension of M to a bicharacter and to an anti-bicharacter gives rise to the same map. Since M is a YB operator, the resulting cochain ψ is effectively a 2-cocycle.

3.1.2. Symmetric twisted tensor products. Given a couple of quantum spaces \mathcal{A} and \mathcal{B} , we recall that a symmetric twisted tensor product (STTP) of them is (essentially) a quantum space $\mathcal{A} \circ_{\tau} \mathcal{B} = (\mathbf{A}_1 \otimes \mathbf{B}_1, \mathbf{A} \circ_{\tau} \mathbf{B})$, being $\mathbf{A} \circ_{\tau} \mathbf{B}$ the subalgebra of $\mathbf{A} \otimes_{\tau} \mathbf{B}$ [1] generated by $\mathbf{A}_1 \otimes \mathbf{B}_1$, and where the related (symmetric) twisting map τ is a linear bijection $\mathbf{B} \otimes \mathbf{A} \simeq \mathbf{A} \otimes \mathbf{B}$ that defines, by restriction, an isomorphism $\mathbf{B}_1 \otimes \mathbf{A}_1 \simeq \mathbf{A}_1 \otimes \mathbf{B}_1$. Thus, τ is completely defined by the last isomorphism and by the properties of a twisting map, namely

$$(3.8) \quad \begin{aligned} \tau(m_B \otimes I_A) &= (I_A \otimes m_B) (\tau \otimes I_B) (I_B \otimes \tau), \\ \tau(I_B \otimes m_A) &= (m_A \otimes I_B) (I_A \otimes \tau) (\tau \otimes I_A), \end{aligned}$$

and

$$(3.9) \quad \tau(I_B \otimes \eta_A) = \eta_A \otimes I_B; \quad \tau(\eta_B \otimes I_A) = I_A \otimes \eta_B.$$

Remember that the algebra structure of $\mathbf{A} \otimes_{\tau} \mathbf{B}$ is given by the maps

$$m_{\mathbf{A} \otimes_{\tau} \mathbf{B}}^{\tau} = (m_A \otimes m_B) (I_A \otimes \tau \otimes I_B) \quad \text{and} \quad \eta_{\mathbf{A} \otimes_{\tau} \mathbf{B}} = \eta_A \otimes \eta_B.$$

We will show that any STTP can be seen as a twisting of $\mathcal{A} \circ \mathcal{B}$ by an element of $\mathfrak{Z}^2[\mathbf{A}_1 \otimes \mathbf{B}_1]$.

The usual product in $\mathbf{A} \otimes \mathbf{B}$ is $m_{\mathbf{A} \otimes \mathbf{B}} = (m_{\mathbf{A}} \otimes m_{\mathbf{B}}) (I \otimes f \otimes I)$, being f the canonical flipping map. Then, given τ we have that $m_{\mathbf{A} \otimes \mathbf{B}}^\tau = m_{\mathbf{A} \otimes \mathbf{B}} \Omega$, where

$$\Omega \doteq (I \otimes f^{-1} \tau \otimes I) : (\mathbf{A} \otimes \mathbf{B})^{\otimes 2} \rightarrow (\mathbf{A} \otimes \mathbf{B})^{\otimes 2}.$$

And since $\Omega \left((\mathbf{A} \circ \mathbf{B})^{\otimes 2} \right) \subset (\mathbf{A} \circ \mathbf{B})^{\otimes 2}$ (which follows from property (3.8)), we also have $m_{\mathbf{A} \circ \mathbf{B}}^\tau = m_{\mathbf{A} \circ \mathbf{B}} \Omega$, by making the corresponding restriction. Now, let us extend the isomorphism

$$\tau|_{\mathbf{B}_1 \otimes \mathbf{A}_1} : \mathbf{B}_1 \otimes \mathbf{A}_1 \simeq \mathbf{A}_1 \otimes \mathbf{B}_1$$

to all of $\mathbf{B}_1^\otimes \otimes \mathbf{A}_1^\otimes$ using Eq. (3.8) and (3.9), and call τ_\otimes the resulting map. By construction, τ_\otimes defines isomorphisms

$$(3.10) \quad \mathbf{B}_1^{\otimes r} \otimes \mathbf{A}_1^{\otimes s} \simeq \mathbf{A}_1^{\otimes s} \otimes \mathbf{B}_1^{\otimes r}, \quad \forall r, s \in \mathbb{N}_0.$$

Then, using the canonical flipping map f_\otimes on $\mathbf{B}_1^\otimes \otimes \mathbf{A}_1^\otimes$ (which obviously defines similar isomorphisms to the ones given in equation above) we have a linear automorphism

$$\widehat{\omega} \doteq (\mathbb{I} \otimes f_\otimes^{-1} \tau_\otimes \otimes \mathbb{I}) : (\mathbf{A}_1^\otimes \otimes \mathbf{B}_1^\otimes)^{\otimes 2} \rightarrow (\mathbf{A}_1^\otimes \otimes \mathbf{B}_1^\otimes)^{\otimes 2}$$

that is homogeneous (since Eq. (3.10)) and that restricted to

$$(\mathbf{A}_1^\otimes \circ \mathbf{B}_1^\otimes)^{\otimes 2} \subset (\mathbf{A}_1^\otimes \otimes \mathbf{B}_1^\otimes)^{\otimes 2}$$

satisfies $\Omega \Pi_\otimes^{\otimes 2} = \Pi_\otimes^{\otimes 2} \widehat{\omega}$. (Recall the identification $\mathbf{A}_1^\otimes \circ \mathbf{B}_1^\otimes \simeq [\mathbf{A}_1 \otimes \mathbf{B}_1]^\otimes$.) Thus, $\widehat{\omega}$ gives rise (by restriction) to a 2-cochain in $\mathcal{C}^2[\mathbf{A}_1 \otimes \mathbf{B}_1]$, which is automatically $\mathcal{A} \circ \mathcal{B}$ -admissible. Let us call ω such a cochain. Because τ_\otimes satisfies Eq. (3.9), ω is counital, i.e. $\omega \in \mathfrak{C}^2[\mathbf{A}_1 \otimes \mathbf{B}_1]$. It remains to see that ω is a 2-cocycle in order to prove our claim. Eq. (3.8) for ω translates into equations $\delta_1 \omega = \omega_{13} \omega_{23}$ and $\delta_2 \omega = \omega_{13} \omega_{12}$, with $\omega_{12} = \omega \otimes \mathbb{I}$, $\omega_{23} = \mathbb{I} \otimes \omega$ and $\omega_{13} = (\mathbb{I} \otimes f_\otimes^{-1}) \omega_{12} (\mathbb{I} \otimes f_\otimes)$. That is to say, ω is an anti-bicharacter. Then, since clearly $\omega_{23} \omega_{12} = \omega_{12} \omega_{23}$, ω is effectively a 2-cocycle; explicitly

$$\delta_1 \omega (\omega \otimes \mathbb{I}) = \omega_{13} \omega_{23} \omega_{12} = \omega_{13} \omega_{12} \omega_{23} = \delta_2 \omega (\mathbb{I} \otimes \omega).$$

Let us resume these results in one proposition.

Proposition 8. *Given an STTP $\mathcal{A} \circ_\tau \mathcal{B}$ of a couple \mathcal{A} and \mathcal{B} of quantum spaces, there exists an admissible counital 2-cocycle ω for $\mathcal{A} \circ \mathcal{B}$ such that $\mathcal{A} \circ_\tau \mathcal{B} = (\mathcal{A} \circ \mathcal{B})_\omega$. Moreover, ω is an anti-bicharacter. ■*

3.1.3. Twisting of bialgebras. Suppose a quantum space \mathcal{A} is a bialgebra in FGA (resp. CA), i.e. the coalgebra structure is given by arrows $\Delta : \mathcal{A} \rightarrow \mathcal{A} \circ \mathcal{A}$ and $\varepsilon : \mathcal{A} \rightarrow \mathcal{I}$ (resp. $\varepsilon : \mathcal{A} \rightarrow \mathcal{K}$) there. In particular, \mathbf{A}_1 is a coalgebra with coproduct $\Delta_1 = \Delta|_{\mathbf{A}_1}$ and counit $\varepsilon_1 = \varepsilon|_{\mathbf{A}_1}$. That is to say, a bialgebra in FGA is a bialgebra algebraically generated by a finite dimensional coalgebra. The pair $(\Delta_1^\otimes, \varepsilon_1^\otimes)$ defines a bialgebra structure on \mathbf{A}_1^\otimes , and the canonical epimorphism Π is a bialgebra homomorphism w.r.t. this structure.

Let $\chi : \mathbf{A}^{\otimes 2} \rightarrow \mathbb{k}$ be a 2-cocycle of $\mathbf{G}^2[\mathbf{A}]$ (see §2.4). From the morphism of quasicomplexes $\mathbf{G}^\bullet[\mathbf{A}] \rightarrow \mathbf{G}^\bullet[\mathbf{A}_1^\otimes]$ induced by the bialgebra map $\Pi : \mathbf{A}_1^\otimes \rightarrow \mathbf{A}$ (see Eq. (2.41) in **Prop. 7**), the map $\chi \Pi^{\otimes 2} : \mathbf{A}_1^\otimes \otimes \mathbf{A}_1^\otimes \rightarrow \mathbb{k}$ is also a 2-cocycle. Since Π is surjective, the maps $\chi \mapsto \chi \Pi^{\otimes n}$, $n \in \mathbb{N}_0$, give rise to a monic $\mathbf{G}^\bullet[\mathbf{A}] \hookrightarrow \mathbf{G}^\bullet[\mathbf{A}_1^\otimes]$. Now, from $F^\bullet : \mathbf{G}^\bullet[\mathbf{A}_1^\otimes] \rightarrow \mathbf{C}^\bullet[\mathbf{A}_1]$, we have in addition the automorphism

$$F_\chi \doteq F^2(\chi \Pi^{\otimes 2}) = (\chi \Pi^{\otimes 2}) * \mathbb{I}^{\otimes 2} * (\chi \Pi^{\otimes 2})^{-1},$$

which, by **Theor. 6**, defines a 2-cocycle in $\mathcal{C}^2[\mathbf{A}_1]$. The composition of F^\bullet with the above monic, quotient out the corresponding centers \mathcal{Z} , is another monic $\mathbf{G}_\mathcal{Z}^\bullet[\mathbf{A}] \hookrightarrow \mathbf{C}^\bullet[\mathbf{A}_1]$ (see **Theor. 7**), which enable us to think of $\mathbf{G}^\bullet[\mathbf{A}]$ as a subcomplex of $\mathbf{C}^\bullet[\mathbf{A}_1]$. Of course, all that is also true if \mathbf{G}^\bullet and \mathbf{C}^\bullet are replaced by their counital subgroups \mathfrak{G}^\bullet and \mathfrak{C}^\bullet .

We recall that the twisting [4][11] of a bialgebra \mathbf{A} by χ is the bialgebra \mathbf{A}_χ with the same coalgebra structure, and associative product $m_\chi = \chi * m * \chi^{-1}$ with the same unit.⁸

F_χ is admissible, in fact

$$(3.11) \quad \Pi^{\otimes 2} F_\chi = \Pi^{\otimes 2} F^2(\chi \Pi^{\otimes 2}) = (\chi * \mathbb{I}^{\otimes 2} * \chi^{-1}) \Pi^{\otimes 2}.$$

Hence, given a counital 2-cocycle $\chi \in \mathbf{G}_\mathcal{Z}^\bullet[\mathbf{A}]$, we have an admissible cocycle F_χ in $\mathfrak{Z}^2[\mathbf{A}_1]$. Moreover, the twisting $\mathcal{A}_\chi = (\mathbf{A}_1, \mathbf{A}_\chi)$ by $\chi \in \mathfrak{G}^2[\mathbf{A}]$ of the bialgebra \mathcal{A} , coincides with the twisting \mathcal{A}_{F_χ} of \mathcal{A} as a quantum space, because from Eq. (3.11), $m_{F_\chi} = \chi * m * \chi^{-1} = m_\chi$.

Let us consider the twisted internal coEnd objects $\text{end}_\mathcal{A}^\Gamma[\mathcal{A}]$, related to a symmetric twisting map defined by $\widehat{\tau}_\mathcal{A} = id \otimes \sigma^\dagger \otimes \sigma$, where $\sigma = \sigma_\mathcal{A} : \mathbf{A}_1 \simeq \mathbf{A}_1$ and $\sigma^\dagger = \sigma^{*-1}$, such that $\sigma_\mathcal{A}$ can be extended to quantum space

⁸If a couple of elements $\chi, \theta \in \mathbf{G}^2$ ‘differ’ by an element of the center \mathcal{Z} , both linear forms define the same twisted bialgebra. This is why the interesting objects are the quotients $\mathbf{G}_\mathcal{Z}^n$ instead of entire groups \mathbf{G}^n .

automorphism $\mathcal{A} \simeq \mathcal{A}$. We mentioned in §1.2 that each one of these bialgebras are isomorphic to the twisting $(\mathcal{A} \triangleright \mathcal{A})_\chi$ (as bialgebras). Such χ is a 2-cocycle in $\mathfrak{G}^2[\mathbf{A} \triangleright \mathbf{A}]$ defined by

$$\chi \left(z_{k_1}^{j_1} \dots z_{k_r}^{j_r} \otimes z_{k_{r+1}}^{j_{r+1}} \dots z_{k_{r+s}}^{j_{r+s}} \right) = \delta_{k_1}^{j_1} \dots \delta_{k_r}^{j_r} (\sigma^{-r})_{k_{r+1}}^{j_{r+1}} \dots (\sigma^{-r})_{k_{r+s}}^{j_{r+s}} \in \mathbb{k},$$

being $z_i^j = a^j \otimes a_i$. Then, via F , $\underline{\text{end}}^\Upsilon[\mathcal{A}]$ can be seen as a twisting of $\underline{\text{end}}[\mathcal{A}]$ by $F_\chi \in \mathfrak{Z}^2[\mathbf{A}_1^* \otimes \mathbf{A}_1]$, where

$$(3.12) \quad \begin{aligned} F_\chi \left(z_{k_1}^{j_1} \dots z_{k_r}^{j_r} \otimes z_{k_{r+1}}^{j_{r+1}} \dots z_{k_{r+s}}^{j_{r+s}} \right) &= \\ &= z_{k_1}^{j_1} \dots z_{k_r}^{j_r} \otimes \left((\sigma^{-r})_{k_{r+1}}^{a_1} \dots (\sigma^{-r})_{k_{r+s}}^{a_s} \right) z_{a_1}^{b_1} \dots z_{a_s}^{b_s} \left((\sigma^r)_{b_1}^{j_{r+1}} \dots (\sigma^r)_{b_s}^{j_{r+s}} \right). \end{aligned}$$

One can show χ , and consequently F_χ , is bicharacter and anti-bicharacter.

In the following we generalize this correspondence between $\underline{\text{end}}^\Upsilon[\mathcal{A}]$ and $\underline{\text{end}}[\mathcal{A}]$ to all of twisted internal coHom objects.

3.1.4. *The twisted coHom objects.* Consider the twisted coHom objects $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ related to a twisting map defined by

$$(3.13) \quad \widehat{\tau}_{\mathcal{A}, \mathcal{B}} = \text{id} \otimes \rho \otimes \phi, \quad \phi = \sigma_{\mathcal{A}}, \quad \rho^{-1} = \sigma_{\mathcal{B}}^*,$$

where $\sigma_{\mathcal{A}}$ and $\sigma_{\mathcal{B}}$ can be extended to automorphisms $\mathcal{A} \simeq \mathcal{A}$ and $\mathcal{B} \simeq \mathcal{B}$. Then, we know that $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ is the algebra generated by $z_i^j = b^j \otimes a_i \in \mathbf{B}_1^* \otimes \mathbf{A}_1$ and quotient by the ideal algebraically generated by (see Eqs. (1.7) to (1.12))

$$\left\{ \left\{ \sigma R_{\lambda_n}^{k_1 \dots k_n} z_{k_1}^{j_1} \cdot z_{k_2}^{j_2} \dots z_{k_n}^{j_n} (\sigma S^\perp)_{j_1 \dots j_n}^{\omega_n} \right\}_{\omega_n \in \Omega_n}^{\lambda_n \in \Lambda_n} \right\}_{n \in \mathbb{N}_0}.$$

Now, let us define for $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] = \mathcal{B} \triangleright \mathcal{A}$ the counital 2-cochain $\varsigma \in \mathfrak{C}^2[\mathbf{B}_1^* \otimes \mathbf{A}_1]$ given by

$$(3.14) \quad \varsigma \left(z_{k_1}^{j_1} \dots z_{k_r}^{j_r} \otimes z_{k_{r+1}}^{j_{r+1}} \dots z_{k_{r+s}}^{j_{r+s}} \right) = z_{k_1}^{j_1} \dots z_{k_r}^{j_r} \otimes \xi^{[r]} \left(z_{k_{r+1}}^{j_{r+1}} \dots z_{k_{r+s}}^{j_{r+s}} \right),$$

being $\xi^{[r]} \in \mathfrak{C}^1[\mathbf{B}_1^* \otimes \mathbf{A}_1]$ such that

$$(3.15) \quad \xi^{[r]} \left(z_{k_1}^{j_1} \dots z_{k_s}^{j_s} \right) = \left((\phi^{-r})_{k_1}^{a_1} \dots (\phi^{-r})_{k_s}^{a_s} \right) z_{a_1}^{b_1} \dots z_{a_s}^{b_s} \left((\rho^{-r})_{b_1}^{j_1} \dots (\rho^{-r})_{b_s}^{j_s} \right).$$

Defining $\xi^{[0]} \doteq \mathbb{I}$, we can write $\varsigma_{r,s} = \mathbb{I}_r \otimes \xi_s^{[r]}$ for $r, s \in \mathbb{N}_0$. From the fact that ϕ and ρ define algebra automorphisms, it follows that ς is $\mathcal{B} \triangleright \mathcal{A}$ -admissible. Let us see that ς is a 2-cocycle. Straightforward calculations show that

$$(3.16) \quad \xi_{s+t}^{[r]} \approx \xi_s^{[r]} \otimes \xi_t^{[r]} \quad \text{and} \quad \xi_t^{[r+s]} = \xi_t^{[r]} \xi_t^{[s]} = \xi_t^{[s]} \xi_t^{[r]}.$$

Then, comparing

$$\varsigma_{r+s,t} (\varsigma_{r,s} \otimes \mathbb{I}_t) = \left(\mathbb{I}_{r+s} \otimes \xi_t^{[r+s]} \right) \left(\mathbb{I}_r \otimes \xi_s^{[r]} \otimes \mathbb{I}_t \right) = \mathbb{I}_r \otimes \xi_s^{[r]} \otimes \xi_t^{[r+s]}$$

and

$$\varsigma_{r,s+t} (\mathbb{I}_r \otimes \varsigma_{s,t}) = \left(\mathbb{I}_r \otimes \xi_{s+t}^{[r]} \right) \left(\mathbb{I}_r \otimes \mathbb{I}_s \otimes \xi_t^{[s]} \right) = \mathbb{I}_r \otimes \xi_{s+t}^{[r]} \left(\mathbb{I}_s \otimes \xi_t^{[s]} \right).$$

we have from Eq. (3.16) that $\varsigma_{r+s,t} (\varsigma_{r,s} \otimes \mathbb{I}_t) \approx \varsigma_{r,s+t} (\mathbb{I}_r \otimes \varsigma_{s,t})$, i.e. ς is an admissible element of $\mathfrak{Z}^2[\mathbf{B}_1^* \otimes \mathbf{A}_1]$. Moreover, using again Eq. (3.16), we can see ς is a bicharacter and anti-bicharacter. The twisting of $\mathcal{B} \triangleright \mathcal{A}$ by ς defines on the vector space $\mathbf{B} \triangleright \mathbf{A}$ the product $m_\varsigma = m \Xi$, with $\Xi \Pi^{\otimes 2} = \Pi^{\otimes 2} \varsigma$. Then, the underlying algebra of $(\mathcal{B} \triangleright \mathcal{A})_\varsigma$ is the one generated by $z_i^j = b^j \otimes a_i$ and quotient by the ideal algebraically generated by

$$\left\{ \left\{ \sigma R_{\lambda_n}^{k_1 \dots k_n} z_{k_1}^{j_1} \cdot z_{k_2}^{j_2} \dots z_{k_n}^{j_n} (\sigma S^\perp)_{j_1 \dots j_n}^{\omega_n} \right\}_{\omega_n \in \Omega_n}^{\lambda_n \in \Lambda_n} \right\}_{n \in \mathbb{N}_0},$$

denoting by \cdot_ς the product m_ς , and the isomorphism $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \simeq \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_\varsigma$ follows. Therefore,

Proposition 9. *The quantum spaces $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ and $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$ are related by a twist transformation, i.e.*

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \simeq \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_\varsigma$$

being ς a bicharacter in $\mathfrak{Z}^2[\mathbf{B}_1^* \otimes \mathbf{A}_1]$. ■

For the coEnd objects the equation

$$\underline{end}[\mathcal{A}]_\chi \simeq \underline{end}^\Upsilon[\mathcal{A}] \simeq \underline{end}[\mathcal{A}]_\varsigma$$

holds, provided $\varsigma = F_\chi$ (see Eq. (3.12)). Moreover, \mathcal{A}_χ and \mathcal{B}_χ can alternatively be seen as twisting of the quantum spaces $\mathcal{A} = \underline{hom}[\mathcal{K}, \mathcal{A}]$ and $\mathcal{B} = \underline{hom}[\mathcal{K}, \mathcal{B}]$ by cocycles

$$a_{k_1} \dots a_{k_r} \otimes a_{k_{r+1}} \dots a_{k_{r+s}} \mapsto a_{k_1} \dots a_{k_r} \otimes (\phi^{-r})_{k_{r+1}}^{j_1} \dots (\phi^{-r})_{k_{r+s}}^{j_s} a_{j_1} \dots a_{j_s},$$

$$b_{k_1} \dots b_{k_r} \otimes b_{k_{r+1}} \dots b_{k_{r+s}} \mapsto b_{k_1} \dots b_{k_r} \otimes (\rho^r)_{k_{r+1}}^{j_1} \dots (\rho^r)_{k_{r+s}}^{j_s} b_{j_1} \dots b_{j_s},$$

in $\mathfrak{Z}^2[\mathbf{A}_1]$ and $\mathfrak{Z}^2[\mathbf{B}_1]$, respectively, which we also denote ς . In resume,

$$\underline{hom}^\Upsilon[\mathcal{B}, \mathcal{A}] \simeq \mathcal{B}_\varsigma \triangleright \mathcal{A}_\varsigma = \underline{hom}[\mathcal{B}_\varsigma, \mathcal{A}_\varsigma] \simeq \underline{hom}[\mathcal{B}, \mathcal{A}]_\varsigma,$$

unifying in this way the correspondence between the objects $\underline{hom}^\Upsilon[\mathcal{B}, \mathcal{A}]$ and $\underline{hom}[\mathcal{B}, \mathcal{A}]$ (in particular, $\mathcal{A}^\Upsilon = \underline{hom}^\Upsilon[\mathcal{K}, \mathcal{A}]$ and $\mathcal{A} = \underline{hom}[\mathcal{K}, \mathcal{A}]$), just in terms of twisting of quantum spaces.

3.2. Twistings and the cohomology relation. We have seen that cocycle condition for a 2-cochain insure the associativity of its related deformed product. But, when can it be insured these products define a twisted quantum space isomorphic to the original one, or when two twisted quantum spaces are isomorphic? The following theorem gives the answer to these questions.

Theorem 9. *Given an \mathcal{A} -admissible cochain $\psi \in \mathfrak{Z}^2$, $\mathcal{A} \simeq \mathcal{A}_\psi$ iff $\psi = \partial\theta$, being θ an \mathcal{A} -admissible counital 1-cochain. More generally, consider a pair of \mathcal{A} -admissible counital 2-cocycles φ and ψ . Then, $\mathcal{A}_\varphi \simeq \mathcal{A}_\psi$ iff φ and ψ are cohomologous through an \mathcal{A} -admissible counital 1-cochain θ . In particular, if φ and ψ are related as in Eq. (3.3), then $\varphi \sim_\theta \psi$ through an admissible θ . ■*

We show below one of the implications. The proof of the other will be given at the end of §3.2.2.

Proof. (part 1) Suppose $\psi = \delta_1\theta (\theta \otimes \theta)^{-1} = \partial\theta$ with

$$\psi(\ker \Pi^{\otimes 2}) = \ker \Pi^{\otimes 2} \quad \text{and} \quad \theta(\ker \Pi) = \ker \Pi.$$

(Since $\partial^2\theta = \mathbb{I}^{\otimes 3}$ for all $\theta \in \mathbf{C}^1$, ψ is a 2-cocycle.) In particular ψ defines a linear automorphism $\Psi : \mathbf{A}^{\otimes 2} \rightarrow \mathbf{A}^{\otimes 2}$ such that $\Psi \Pi^{\otimes 2} = \Pi^{\otimes 2} \psi$, which gives rise to the twisted product $m_\psi = m\Psi$. And θ defines, by the equation $\Theta \Pi = \Pi\theta$, a linear automorphism $\Theta : \mathbf{A} \rightarrow \mathbf{A}$ satisfying $\Theta(\mathbf{A}_1) \subset \mathbf{A}_1$. Then,

$$m_\psi \Pi^{\otimes 2} = m \Pi^{\otimes 2} \psi = m \Pi^{\otimes 2} \delta_1\theta (\theta \otimes \theta)^{-1} = \Pi m_\otimes \delta_1\theta (\theta \otimes \theta)^{-1} = \Pi\theta m_\otimes (\theta \otimes \theta)^{-1},$$

where Eq. (2.7) was used in the last equality. Multiplying to the right by $\theta \otimes \theta$ and using the relation between θ and Θ we arrive at $m_\psi (\Theta \otimes \Theta) \Pi^{\otimes 2} = \Theta m \Pi^{\otimes 2}$, and since Π is right invertible, equality $m_\psi (\Theta \otimes \Theta) = \Theta m$ follows. This means the linear automorphism Θ defines an algebra map $\Theta : \mathbf{A} \rightarrow \mathbf{A}_\psi$ such that $\Theta(\mathbf{A}_1) \subset \mathbf{A}_1$, therefore it defines a quantum space isomorphism $\mathcal{A} \simeq \mathcal{A}_\psi$.

Now, consider a pair of admissible 2-cochains φ and ψ such that $\varphi \sim_\theta \psi$, being θ admissible. By definition of cohomologous $\delta_1\theta \varphi (\theta \otimes \theta)^{-1} = \psi$. Hence, reproducing the calculations above, $m_\psi (\Theta \otimes \Theta) = \Theta m_\varphi$, being Θ the linear automorphism defined by θ and Π . That is to say $\mathcal{A}_\varphi \simeq \mathcal{A}_\psi$. ■

Consider on the set of \mathcal{A} -admissible counital 2-cocycles, the cohomology relation through \mathcal{A} -admissible 1-cochains; i.e. the \mathcal{A} -admissible cochains $\varphi, \psi \in \mathfrak{Z}^2$ are cohomologous if $\varphi \sim_\theta \psi$, being θ admissible. It is an equivalence relation since proposition of §2.2.3 and the fact that admissible n -cochains form a subgroup of $\mathbf{C}^n[\mathbf{A}_1]$ for all n . Then, by theorem above, the resulting quotient set, namely $H_{\mathcal{A}}^2[\mathbf{A}_1]$, characterize completely the isomorphism classes of twisting of \mathcal{A} . Since the last claim of theorem above (related to Eq. (3.3)), the set $H_{\mathcal{A}}^2[\mathbf{A}_1]$ can be regarded as a quotient of $Aut_{\text{Vct}}[\mathbf{A} \otimes \mathbf{A}]$.

Following an analogous reasoning, one can define a space $H_{\mathcal{A}}^1[\mathbf{A}_1]$, given by \mathcal{A} -admissible 1-cocycles (*ipso facto* counital) quotient by the cohomology relation through admissible cochains (which gives the equality relation, as follows from the general case analyzed in §2.2.1). This space is in bijection with the group of automorphisms of \mathcal{A} , i.e. $H_{\mathcal{A}}^1[\mathbf{A}_1] \simeq Aut_{\text{FGA}}[\mathcal{A}]$.⁹

It is worth mentioning that **non** every admissible 2-cocycle is a coboundary of an admissible 1-cochain. If this were the case, the commutative algebra $\mathbb{k}[a, b]$ of §3.1.1 would be isomorphic to the quantum plane $\mathbf{A}_\hbar^{2|0} = \mathbb{k}[a, b]_{\psi_\hbar}$. In particular, we can affirm that there do exist non trivial twisting of quantum spaces, i.e. $H_{\mathcal{A}}^2[\mathbf{A}_1] \neq \{\mathbb{I}^{\otimes 2}\}$. Nevertheless, for quantum spaces formed out by free algebras, every counital 2-cocycle gives

⁹To give an automorphism of $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ is the same as to give an element $\alpha \in Aut[\mathbf{A}_1]$ such that α, α^{-1} can be extended to algebra maps on \mathbf{A} . But these are precisely the admissible 1-cocycles in $\mathbf{C}^1[\mathbf{A}_1]$. Since $\mathbf{Z}^1[\mathbf{A}_1] = \mathfrak{Z}^1[\mathbf{A}_1] \simeq Aut[\mathbf{A}_1]$, we have the mentioned bijection.

rise to an isomorphic twisted quantum space.¹⁰ This is a consequence of the fact that every counital 2-cocycle is a counital 2-coboundary (and the admissibility condition is immediate), as we show below.

3.2.1. *The equality $Z^2 = B^2$.* We have seen that $B^2 \subset Z^2$ for every vector space. In this subsection we show the other inclusion and some related consequences.

Theorem 10. *Every $\psi \in Z^2$ is cohomologous to the identity, i.e. $\psi \in B^2$. Moreover, for every linear automorphism $\varpi : \mathbf{A}_1 \simeq \mathbf{A}_1$, there exists a unique $\theta \in C^1$ such that $\psi = \partial\theta$ and $\theta_1 = \varpi$.*

Proof. Let us come back to the proof of **Theor. 8**. Given a cochain $\psi \in C^2$ (non necessarily a counital 2-cocycle) we have defined inductively, from Eq. (3.5), a family of linear isomorphisms $\theta_n : \mathbf{A}_1^{\otimes n} \simeq \mathbf{A}_1^{\otimes n}$, $n \geq 2$. In an analogous way, take a map $\varpi : \mathbf{A}_1 \simeq \mathbf{A}_1$ and consider $\theta \in C^1$ such that $\theta_0 \doteq \psi_{0,0}^{-1} = 1/\psi_{0,0}$, $\theta_1 \doteq \varpi$, and

$$\theta_{n+1} \doteq \psi_{n,1}(\theta_n \otimes \varpi), \quad n \in \mathbb{N}.$$

Let us show that if ψ is a 2-cocycle then, using the identification given by Eq. (2.8),

$$(3.17) \quad \theta_{r+s} \approx \psi_{r,s}(\theta_r \otimes \theta_s),$$

or equivalently $\psi = \delta_1\theta(\theta \otimes \theta)^{-1} = \partial\theta$. We will make induction on s . Recall that, since Eq. (2.20), $\psi_{r,0} \approx \psi_{0,0} \cdot \mathbb{I}_r$ for all 2-cocycle. Then for $s = 0$, by definition of θ_0 and θ_1 , Eq. (3.17) holds. For $s + 1$,

$$\psi_{r,s+1}(\theta_r \otimes \theta_{s+1}) \approx \psi_{r,s+1}(\theta_r \otimes \psi_{s,1}(\theta_s \otimes \varpi)) = \psi_{r,s+1}(\mathbb{I}_r \otimes \psi_{s,1})(\theta_r \otimes \theta_s \otimes \varpi).$$

Using Eq. (2.19) for $t = 1$,

$$\psi_{r,s+1}(\theta_r \otimes \theta_{s+1}) \approx \psi_{r+s,1}(\psi_{r,s} \otimes \mathbb{I})(\theta_r \otimes \theta_s \otimes \varpi) = \psi_{r+s,1}(\psi_{r,s}(\theta_r \otimes \theta_s) \otimes \varpi),$$

and from the inductive hypothesis (i.e. validity of Eq. (3.17))

$$\psi_{r,s+1}(\theta_r \otimes \theta_{s+1}) \approx \psi_{r+s,1}(\theta_{r+s} \otimes \varpi) = \theta_{r+s+1},$$

as we wanted to show.

Now, consider another χ such that $\psi = \partial\chi$ and $\chi_1 = \varpi$. From (3.17), $\chi_0 = 1/\psi_{0,0}$ and the rest of χ is given inductively by the equation $\chi_{n+1} = \psi_{n,1}(\chi_n \otimes \varpi)$, concluding in this way the proof of the our claim. ■

Corollary 1. *If $\psi \in \mathfrak{Z}^2$, then $\psi \in \mathfrak{B}^2$. In addition, fixing an automorphism ϖ of \mathbf{A}_1 , there exists a unique $\theta \in C^1$ such that $\psi = \partial\theta$ and $\theta_1 = \varpi$.*

Proof. From previous theorem, if $\psi \in \mathfrak{Z}^2 = Z^2 \cap \mathfrak{C}^2$, then $\psi \in B^2 \cap \mathfrak{C}^2$ and there exists a unique $\theta \in C^1$ such that $\psi = \partial\theta$ and $\theta_1 = \varpi$. But, as we have shown in §2.2.1 (see Eq. (2.22)), $B^2 \cap \mathfrak{C}^2 = \mathfrak{B}^2$. Thus $\psi \in \mathfrak{B}^2$ and accordingly $\theta \in C^1$. ■

This corollary can also be proven by noting that ψ is counital if and only if $\psi_{0,0} = 1$ (see Eq. (2.21)). This makes $\theta_0 = \mathbb{I}_0 = 1$, i.e. θ is counital.

Corollary 2. *Every twisting of the quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A}_1^{\otimes})$ is isomorphic to \mathcal{A} .*

Proof. Every n -cochain in $C^n[\mathbf{A}_1]$ is obviously admissible for the quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A}_1^{\otimes})$ (since $\ker \Pi = 0$). Then, from above results, given a counital 2-cocycle ψ there exists a counital 1-cochain θ (admissible for \mathcal{A}) such that $\psi = \partial\theta$. In this situation, **Theor. 9** insures $\mathcal{A} \simeq \mathcal{A}_\psi$. ■

These results motive the definition of a subgroup $\mathfrak{P}^1[\mathbf{A}_1] \subset C^1[\mathbf{A}_1]$ constituted by counital 1-cochains θ such that $\theta_1 = \mathbb{I}$. We shall call **primitive** the elements of this subgroup. From now on, when we write $\psi = \partial\theta$, we are supposing θ is in \mathfrak{P}^1 , unless we say the contrary.

As an example, the cochain

$$(3.18) \quad \varsigma_{\mathcal{A}} : a_{k_1} \dots a_{k_r} \otimes a_{k_{r+1}} \dots a_{k_{r+s}} \mapsto a_{k_1} \dots a_{k_r} \otimes (\sigma_{\mathcal{A}}^{-r})_{k_{r+1}}^{j_1} \dots (\sigma_{\mathcal{A}}^{-r})_{k_{r+s}}^{j_s} a_{j_1} \dots a_{j_s}$$

related to the twisted internal coHom objects $\mathcal{B}_\varsigma \triangleright \mathcal{A}_\varsigma$, has primitive

$$\theta_{\mathcal{A}} : a_{k_1} \dots a_{k_r} \mapsto \delta_{k_1}^{j_1} (\sigma_{\mathcal{A}}^{-1})_{k_2}^{j_2} (\sigma_{\mathcal{A}}^{-2})_{k_3}^{j_3} \dots (\sigma_{\mathcal{A}}^{-r+1})_{k_r}^{j_r} a_{j_1} \dots a_{j_r}.$$

Since we have shown in §2.2.3 that cohomology relation is an equivalence relation in $C^2[\mathbf{A}_1]$, previous results imply, in particular, that every pair of 2-cocycles are cohomologous. The following proposition tell us which class of 1-cochains implement such a relation.

¹⁰If the twisted space is not also free, then it would have a non zero related ideal, which would imply the underlying vector space is smaller than the untwisted one.

Proposition 10. Consider a pair of cochains $\psi, \varphi \in \mathfrak{Z}^2$ with primitive $\lambda, \chi \in \mathfrak{P}^1$, respectively.

a) They are cohomologous through θ iff $\theta = \lambda \omega \chi^{-1}$, with $\omega \in \mathfrak{Z}^1$.

b) They are cohomologous through an admissible cochain iff there exists $\omega \in \mathfrak{Z}^1$ such that $\lambda \omega \chi^{-1}$ is admissible.

Proof. a) If $\psi = \partial\lambda$, $\varphi = \partial\chi$, $\theta = \lambda \omega \chi^{-1}$ and $\partial\omega = \mathbb{I}^{\otimes 2}$ (i.e. ω is a 1-cocycle), then

$$\begin{aligned} \delta_1 \theta \varphi (\theta \otimes \theta)^{-1} &= \delta_1 (\lambda \omega \chi^{-1}) \left(\delta_1 \chi (\chi \otimes \chi)^{-1} \right) (\lambda \omega \chi^{-1} \otimes \lambda \omega \chi^{-1})^{-1} \\ &= \delta_1 (\lambda \omega) (\lambda \omega \otimes \lambda \omega)^{-1} = \delta_1 \lambda \left(\delta_1 \omega (\omega \otimes \omega)^{-1} \right) (\lambda \otimes \lambda)^{-1} = \delta_1 \lambda (\lambda \otimes \lambda)^{-1} = \psi. \end{aligned}$$

That means $\varphi \smile_{\theta} \psi$, and one implication follows. Now suppose $\varphi \smile_{\theta} \psi$. We always can write $\theta = \lambda \omega \chi^{-1}$, for every $\theta \in \mathbf{C}^1$. It is enough to take $\omega = \lambda^{-1} \theta \chi$. Let us see that $\omega \in \mathfrak{Z}^1$. From equation above if $\varphi \smile_{\theta} \psi$ we have that

$$\delta_1 \lambda \left(\delta_1 \omega (\omega \otimes \omega)^{-1} \right) (\lambda \otimes \lambda)^{-1} = \delta_1 \lambda (\lambda \otimes \lambda)^{-1},$$

which is fulfilled if and only if $\delta_1 \omega = \omega \otimes \omega$, i.e. $\omega \in \mathfrak{Z}^1$, as we wanted to show.

b) This part is an immediate consequence of the first one. ■

Recall that to give an element $\omega \in \mathfrak{Z}^1[\mathbf{A}_1]$ is the same as giving one in $\varpi \in \text{Aut}[\mathbf{A}_1]$, since every 1-cocycle is of the form $\omega = \varpi^{\otimes}$. This connects above proposition with **Theor. 9**.

A brief comment about bialgebras is in order. The quasicomplex $\mathbf{G}^{\bullet}[\mathbf{A}_1^{\otimes}]$ also satisfies that every 2-cocycle is a 2-coboundary. In fact, given a cocycle $\chi \in \mathbf{G}^2$, a cochain $\lambda \in \mathbf{G}^1$, such that $d\lambda = \chi$, can be inductively defined by the formula

$$\lambda_0 = 1/\chi_{0,0}; \quad \lambda_{n+1} = (\lambda_n \otimes \varepsilon) * \chi_{n,1}.$$

Moreover, every pair of 2-cocycles in $\mathbf{G}_{\mathbb{Z}}^2[\mathbf{A}_1^{\otimes}]$ are cohomologous.¹¹ But for $\mathbf{G}_{\mathbb{Z}}^{\bullet}[\mathbf{A}]$, given a 2-cocycle ϕ there, we can just say $\phi \Pi^{\otimes 2} = d\vartheta$; and ϕ will be a coboundary iff $\vartheta = \theta \Pi$.

3.2.2. Role of primitive 1-cochains. The role of the 1-cochains θ defining a given 2-cocycle ψ , for a generic quantum space \mathcal{A} , is expressed by the following theorem.

Theorem 11. Let $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ be a quantum space and consider an admissible $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$. The canonical epimorphism $\mathbf{A}_1^{\otimes} \rightarrow \mathbf{A}_{\psi}$ associated to $\mathcal{A}_{\psi} = (\mathbf{A}_1, \mathbf{A}_{\psi})$ is given by $\Pi\theta$, being $\theta \in \mathfrak{P}^1[\mathbf{A}_1]$ the primitive of ψ . Furthermore, if \mathbf{I} is the ideal related to \mathcal{A} , then \mathbf{I}_{ψ} , the one related to \mathcal{A}_{ψ} , is equal to $\theta^{-1}(\mathbf{I})$.

Proof. Since $\theta_{0,1} = \mathbb{I}_{0,1}$ (which means $\theta(\mathbf{A}_1) \equiv \mathbf{A}_1$ and $\theta(\lambda) = \eta(\lambda)$), $\Pi\theta$ defines the inclusion $\mathbf{A}_1 \hookrightarrow \mathbf{A}_{\psi}$ and a unit preserving map. If we show that

$$(3.19) \quad m_{\otimes} \psi (\theta \otimes \theta) = \theta m_{\otimes},$$

what implies (multiplying by Π to the left) $m_{\psi} (\Pi\theta \otimes \Pi\theta) = \Pi\theta m_{\otimes}$, we are proving $\Pi\theta$ is an algebra homomorphism $\mathbf{A}_1^{\otimes} \rightarrow \mathbf{A}_{\psi}$.

Eq. (3.19) restricted to a subspace $\mathbf{A}_1^{\otimes r} \otimes \mathbf{A}_1^{\otimes s}$ means $\psi_{r,s} (\theta_r \otimes \theta_s) = [\delta_1 \theta]_{r,s}$, or equivalently, $\psi_{r,s} (\theta_r \otimes \theta_s) \approx \theta_{r+s}$. But this is true since $\partial\theta = \psi$, hence Eq. (3.19) holds.

Finally, we must show $\theta^{-1}(\mathbf{I})$ is equal to \mathbf{I}_{ψ} , the ideal related to \mathcal{A}_{ψ} . By definition, $\mathbf{I} = \ker \Pi$ and, from the last result, $\mathbf{I}_{\psi} = \ker \Pi\theta$. In addition, it is well known that $\ker \Pi\theta = \theta^{-1}(\ker \Pi)$, and consequently our claim follows. ■

Thus we can write (compare to Eq. (3.4))

$$a_{j_1} \cdot_{\psi} \dots \cdot_{\psi} a_{j_n} = \Pi\theta (a_{i_1} \otimes \dots \otimes a_{i_n}),$$

being " \cdot_{ψ} " = m_{ψ} . On the other hand, note that if θ is \mathcal{A} -admissible, then $\mathbf{I}_{\psi} = \theta^{-1}(\mathbf{I}) = \mathbf{I}$ and, therefore, $\mathcal{A} \simeq \mathcal{A}_{\psi}$, as we proved before.

As an immediate corollary of last theorem we have:

Corollary 3. Consider a conic quantum space \mathcal{A} with $\mathbf{I} = \bigoplus_{n \geq 2} \mathbf{I}_n$, and a counital \mathcal{A} -admissible 2-cocycle $\psi = \partial\theta \in \mathfrak{Z}^2[\mathbf{A}_1]$. The ideal related to the twisted quantum space \mathcal{A}_{ψ} is

$$\mathbf{I}_{\psi} = \bigoplus_{n \geq 2} \mathbf{I}_{\psi,n} = \bigoplus_{n \geq 2} \theta^{-1}(\mathbf{I}_n). \quad \blacksquare$$

¹¹The cochains that implement the cohomology relation have a form analogous to the ones given in last proposition for the complex $\mathbf{C}^{\bullet}[\mathbf{A}_1]$.

In coordinates, if \mathbf{I} is linearly generated by

$$\left\{ R_{\lambda_n}^{k_1 \dots k_n} a_{k_1 \dots k_n} \right\}_{\lambda_n \in \Lambda_n} \subset \mathbf{I}_n,$$

then \mathbf{I}_ψ is linearly generated by the elements

$$R_{\lambda_n}^{k_1 \dots k_n} \theta^{-1}(a_{k_1 \dots k_n}) = R_{\lambda_n}^{k_1 \dots k_n} (\theta_n^{-1})_{k_1 \dots k_n}^{j_1 \dots j_n} a_{j_1 \dots j_n}.$$

Last corollary gives us another way to see that twisting of conic quantum spaces are also conic. Further work shows the same for m -th quantum spaces. In what follows, and to avoid any confusion, we shall write $\mathbf{X} \cdot \mathbf{Y} \subset \mathbf{A}_1^\otimes$ for the image under m_\otimes of a couple of subspaces $\mathbf{X}, \mathbf{Y} \subset \mathbf{A}_1^\otimes$.

Proposition 11. *If $\mathcal{A} \in \text{CA}^m$ and $\mathbf{I} = I[\mathbf{Y}]$, the ideal generated by $\mathbf{Y} \subset \mathbf{A}_1^{\otimes m}$, then $\mathcal{A}_\psi \in \text{CA}^m$ and $\mathbf{I}_\psi = I[\theta^{-1}(\mathbf{Y})]$. That is to say, twist transformations of $\mathcal{A} \in \text{CA}^m$ give objects of CA^m . Moreover, if $\mathcal{A} \in \text{CA}$ has an ideal \mathbf{I} generated by a graded vector subspace of \mathbf{A}_1^\otimes*

$$\mathbf{S} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{S}_n; \quad \mathbf{S}_{0,1} \doteq \{0\},$$

i.e. $\mathbf{I} = I[\mathbf{S}] = \mathbf{A}_1^\otimes \cdot \mathbf{S} \cdot \mathbf{A}_1^\otimes$,¹² then the ideal related to \mathcal{A}_ψ is

$$\mathbf{I}_\psi = \theta^{-1}(I[\mathbf{S}]) = I[\theta^{-1}(\mathbf{S})] = \mathbf{A}_1^\otimes \cdot \theta^{-1}(\mathbf{S}) \cdot \mathbf{A}_1^\otimes. \quad \blacksquare$$

Before going to the proof, let us show the lemma below.

Lemma 3. *Let \mathcal{A} be a conic quantum space with ideal $\mathbf{I} = \bigoplus_{n \geq 2} \mathbf{I}_n$. $\psi = \partial\theta$ is admissible iff*

$$(3.20) \quad \theta^{-1}(\mathbf{A}_1^{\otimes r} \cdot \mathbf{I}_s + \mathbf{I}_r \cdot \mathbf{A}_1^{\otimes s}) = \mathbf{A}_1^{\otimes r} \cdot \theta^{-1}(\mathbf{I}_s) + \theta^{-1}(\mathbf{I}_r) \cdot \mathbf{A}_1^{\otimes s}.$$

Proof. If ψ is admissible, then (see Eq. (3.2))

$$\psi_{r,s}(\mathbf{A}_1^{\otimes r} \otimes \mathbf{I}_s + \mathbf{I}_r \otimes \mathbf{A}_1^{\otimes s}) = \mathbf{A}_1^{\otimes r} \otimes \mathbf{I}_s + \mathbf{I}_r \otimes \mathbf{A}_1^{\otimes s}.$$

Since $\psi = \delta_1\theta (\theta \otimes \theta)^{-1}$, equation above says

$$(3.21) \quad [\delta_1\theta^{-1}]_{r,s}(\mathbf{A}_1^{\otimes r} \otimes \mathbf{I}_s + \mathbf{I}_r \otimes \mathbf{A}_1^{\otimes s}) = (\theta_r \otimes \theta_s)^{-1}(\mathbf{A}_1^{\otimes r} \otimes \mathbf{I}_s + \mathbf{I}_r \otimes \mathbf{A}_1^{\otimes s}).$$

Applying to the left m_\otimes , using $m_\otimes \delta_1\theta = \theta m_\otimes$ and $\theta(\mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes r}$, we arrive precisely at Eq. (3.20). Reciprocally, if θ satisfies (3.20), Eq. (3.21) follows immediately and with it the admissibility of $\partial\theta$. \blacksquare

Proof. (of proposition) Let us consider the more general case. Suppose we have $\mathcal{A} \in \text{CA}$ with

$$I[\mathbf{S}] = \bigoplus_{n \geq 2} \mathbf{I}_n; \quad \mathbf{I}_n = \sum_{r=2}^n \sum_{i=0}^{n-r} \mathbf{A}_1^{\otimes n-r-i} \cdot \mathbf{S}_r \cdot \mathbf{A}_1^{\otimes i},$$

and an \mathcal{A} -admissible 2-cocycle $\psi = \partial\theta$. From lemma above, since $\mathbf{I}_1 = \{0\}$, we have

$$(3.22) \quad \theta^{-1}(\mathbf{A}_1 \cdot \mathbf{I}_r) = \mathbf{A}_1 \cdot \theta^{-1}(\mathbf{I}_r) \quad \text{and} \quad \theta^{-1}(\mathbf{I}_r \cdot \mathbf{A}_1) = \theta^{-1}(\mathbf{I}_r) \cdot \mathbf{A}_1,$$

for all r , and in particular

$$\theta^{-1}(\mathbf{A}_1 \cdot \mathbf{S}_2) = \mathbf{A}_1 \cdot \theta^{-1}(\mathbf{S}_2) \quad \text{and} \quad \theta^{-1}(\mathbf{S}_2 \cdot \mathbf{A}_1) = \theta^{-1}(\mathbf{S}_2) \cdot \mathbf{A}_1.$$

Consequently, $\theta^{-1}(\mathbf{I}_3) = \mathbf{A}_1 \cdot \theta^{-1}(\mathbf{S}_2) + \theta^{-1}(\mathbf{S}_2) \cdot \mathbf{A}_1 + \theta^{-1}(\mathbf{S}_3)$. We can show by induction on $n \geq 3$ that

$$\theta^{-1}(\mathbf{I}_n) = \sum_{r=2}^n \sum_{i=0}^{n-r} \mathbf{A}_1^{\otimes n-r-i} \cdot \theta^{-1}(\mathbf{S}_r) \cdot \mathbf{A}_1^{\otimes i}.$$

In fact, because $\mathbf{I}_{n+1} = \mathbf{A}_1 \cdot \mathbf{I}_n + \mathbf{I}_n \cdot \mathbf{A}_1 + \mathbf{S}_{n+1}$, and using (3.22)

$$\theta^{-1}(\mathbf{I}_{n+1}) = \theta^{-1}(\mathbf{A}_1 \cdot \mathbf{I}_n + \mathbf{I}_n \cdot \mathbf{A}_1 + \mathbf{S}_{n+1}) = \mathbf{A}_1 \cdot \theta^{-1}(\mathbf{I}_n) + \theta^{-1}(\mathbf{I}_n) \cdot \mathbf{A}_1 + \theta^{-1}(\mathbf{S}_{n+1}),$$

we have from inductive hypothesis

$$\begin{aligned} \theta^{-1}(\mathbf{I}_{n+1}) &= \sum_{r=2}^n \sum_{i=0}^{n-r} \mathbf{A}_1^{\otimes n+1-r-i} \cdot \theta^{-1}(\mathbf{S}_r) \cdot \mathbf{A}_1^{\otimes i} \\ &+ \sum_{r=2}^n \sum_{i=0}^{n-r} \mathbf{A}_1^{\otimes n-r-i} \cdot \theta^{-1}(\mathbf{S}_r) \cdot \mathbf{A}_1^{\otimes i+1} + \theta^{-1}(\mathbf{S}_{n+1}) = \sum_{r=2}^{n+1} \sum_{i=0}^{n+1-r} \mathbf{A}_1^{\otimes n+1-r-i} \cdot \theta^{-1}(\mathbf{S}_r) \cdot \mathbf{A}_1^{\otimes i}. \end{aligned}$$

Thus, $\theta^{-1}(I[\mathbf{S}]) = I[\theta^{-1}(\mathbf{S})]$, as we wanted to see. \blacksquare

¹²This is the case of quantum spaces of the form $\mathcal{A} \circ \mathcal{B}$ with $\circ = \bullet, \triangleleft, \triangleright, \diamond$ or \odot , and $\mathcal{A}^!$.

Example: In the quadratic case $\mathbf{I} = I[\mathbf{Y}]$ is an ideal generated by $\mathbf{Y} \subset \mathbf{A}_1^{\otimes 2}$. Then, since $\theta_2 \doteq \psi_{1,1}$, we have that $\mathbf{I}_\psi = I[\psi_{1,1}^{-1}(Y)]$. Let us suppose \mathbf{Y} is given by a set $Y_\lambda^{kl} a_k \otimes a_l$, $\lambda \in \Lambda$. If $\psi_{1,1} \doteq M \in GL(n^2)$, \mathbf{I}_ψ is algebraically generated by the elements $Y_\lambda^{kl} N_{kl}^{ij} a_i \otimes a_j$, with $N = M^{-1}$. For a freely commutative algebra, we have $\Lambda = [n] \times [n]$ and $Y_\lambda^{kl} = Y_{ij}^{kl} = \delta_i^k \delta_j^l - P_{ij}^{kl}$. From all that Equation (3.7) follows. ■

To end this subsection, let us prove the other implication of **Theor. 9**.

Proof. (of Theor. 9, part 2) Suppose $\mathcal{A}_\varphi \simeq \mathcal{A}_\psi$. That is to say, there exists a linear automorphism $\Theta : \mathbf{A} \simeq \mathbf{A}$ such that $\Theta(1) = 1$ and, in a basis $\{a_i\}$ of \mathbf{A}_1 , $\Theta(a_i) = \varpi_i^j a_j$ and

$$\Theta(a_{i_1} \cdot_\varphi \dots \cdot_\varphi a_{i_k}) = \varpi_{i_1}^{j_1} \dots \varpi_{i_k}^{j_k} a_{j_1} \cdot_\psi \dots \cdot_\psi a_{j_k},$$

being ϖ_i^j the matrix elements of an automorphism $\varpi : \mathbf{A}_1 \simeq \mathbf{A}_1$. If χ and λ are the primitive of φ and ψ , resp., the last equation can be written $\Theta \Pi \chi_k = \Pi \lambda_k \varpi^{\otimes k}$, $k \in \mathbb{N}_0$. Thus, the 1-cochain θ given by $\theta_k = \lambda_k \varpi^{\otimes k} \chi_k^{-1}$ is \mathcal{A} -admissible. Calling ω the 1-cocycle such that $\omega_r = \varpi^{\otimes r}$, then $\theta = \lambda \omega \chi^{-1}$. From **Prop. 10** it follows that $\varphi \sim_\theta \psi$. In particular, if $\Pi^{\otimes 2}(\varphi - \psi) = 0$ (see Eq. (3.3)), then $m_\varphi = m_\psi$ and, in consequence, $\mathcal{A}_\varphi = \mathcal{A}_\psi$. That implies φ and ψ are cohomologous through the admissible 1-cochain $\lambda \chi^{-1}$ (since $\varpi = \mathbb{I}$). ■

3.3. The gauge equivalence.

3.3.1. *Composition and inversion of twist transformations.* In this subsection we show that consecutive applications of twist transformations is again a twist transformation, and that twist transformations have inverse. To start with, the following results will be crucial.

Proposition 12. *The map ∂ defines a surjection $\mathfrak{C}^1 \rightarrow \mathfrak{Z}^2$ which restricted to the subgroup $\mathfrak{P}^1 \subset \mathfrak{C}^1$ became a bijection $\mathfrak{P}^1 \simeq \mathfrak{Z}^2$. ■*

This is an immediate corollary of **Theor. 10**.

Proposition 13. *The set \mathfrak{Z}^2 is a group under the product*

$$(\psi, \varphi) \mapsto \psi \star \varphi \doteq \psi(\theta \otimes \theta) \varphi(\theta \otimes \theta)^{-1},$$

being $\theta \in \mathfrak{P}^1$ the primitive of ψ . With respect to this group structure the bijection $\mathfrak{P}^1 \simeq \mathfrak{Z}^2$ is a group isomorphism.

Proof. From $\partial : \mathfrak{P}^1 \simeq \mathfrak{Z}^2$ we have that $\psi \star \varphi = \partial\theta \star \partial\chi$ is equal to

$$\partial\theta(\theta \otimes \theta) \partial\chi(\theta \otimes \theta)^{-1} = \delta_1 \theta \delta_1 \chi(\theta \otimes \theta)^{-1} (\chi \otimes \chi)^{-1} = \delta_1 \theta (\theta \chi \otimes \theta \chi)^{-1} = \partial(\theta \chi),$$

thus $\theta \chi \in \mathfrak{P}^1$ is sent to $\psi \star \varphi$ via ∂ . In other words, ∂ restricted to \mathfrak{P}^1 translates the product on its domain into the map \star . Hence, \star is an associative product with unit $\mathbb{I}^{\otimes 2} = \partial\mathbb{I}$, and each $\psi \in \mathfrak{Z}^2$ has inverse

$$i\psi = \partial(\theta^{-1}) = (\delta_1 \theta)^{-1} (\theta \otimes \theta) = (\theta \otimes \theta)^{-1} (\partial\theta)^{-1} (\theta \otimes \theta) = (\theta \otimes \theta)^{-1} \psi^{-1} (\theta \otimes \theta).$$

This concludes the proof. ■

Now we are ready to prove our claim.

Proposition 14. *Given a quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ and a couple of counital 2-cocycles ψ and φ , admissible for \mathcal{A} and \mathcal{A}_ψ , respectively, then $\psi \star \varphi$ is \mathcal{A} -admissible and $(\mathcal{A}_\psi)_\varphi = \mathcal{A}_{\psi \star \varphi}$. In addition $i\psi$, the inverse of ψ under the product \star , is admissible for \mathcal{A}_ψ , and accordingly $(\mathcal{A}_\psi)_{i\psi} = \mathcal{A}$.*

Proof. Let Ψ and Φ be the automorphisms of $\mathbf{A}^{\otimes 2}$ related to ψ and φ , respectively, i.e.

$$\Psi \Pi^{\otimes 2} = \Pi^{\otimes 2} \psi \quad \text{and} \quad \Phi (\Pi \theta)^{\otimes 2} = (\Pi \theta)^{\otimes 2} \varphi,$$

if $\psi = \partial\theta$. The last equality implies $\Phi \Pi^{\otimes 2} = \Pi^{\otimes 2} (\theta \otimes \theta) \varphi (\theta \otimes \theta)^{-1}$ and

$$\Psi \Phi \Pi^{\otimes 2} = \Pi^{\otimes 2} \psi (\theta \otimes \theta) \varphi (\theta \otimes \theta)^{-1} = \Pi^{\otimes 2} \psi \star \varphi.$$

Thus $\psi \star \varphi$ is \mathcal{A} -admissible, and since the product of $(\mathcal{A}_\psi)_\varphi$ is $(m\Psi)\Phi = m(\Psi\Phi)$, the first claim of the proposition follows. Now, we must show that the 2-cocycle $i\psi = \partial(\theta^{-1}) = (\theta \otimes \theta)^{-1} \psi^{-1} (\theta \otimes \theta)$ is admissible for \mathcal{A}_ψ . But

$$(3.23) \quad (\Pi \theta)^{\otimes 2} i\psi = \Pi^{\otimes 2} \psi^{-1} (\theta \otimes \theta) = \Psi^{-1} (\Pi \theta)^{\otimes 2},$$

hence $i\psi \left(\ker(\Pi \theta)^{\otimes 2} \right) \subset \ker(\Pi \theta)^{\otimes 2}$. Then, the twisting of \mathcal{A}_ψ by $i\psi$ can be defined, and from the above result $(\mathcal{A}_\psi)_{i\psi} = \mathcal{A}_{\psi \star i\psi} = \mathcal{A}$. In particular, the multiplication of $(\mathcal{A}_\psi)_{i\psi}$ will be, since Eq. (3.23), $(m\Psi)\Psi^{-1} = m(\Psi\Psi^{-1}) = m$. ■

Let us recall that something similar happens in the bialgebra case. It is well-known that if χ, ζ are 2-cocycles in $G^2[\mathbf{A}]$ and $G^2[\mathbf{A}_\chi]$, respectively, then $\zeta * \chi$ is a 2-cocycle in $G^2[\mathbf{A}]$, and $(\mathbf{A}_\chi)_\zeta = \mathbf{A}_{\zeta * \chi}$. In addition, χ^{-1} is a 2-cocycle in $G^2[\mathbf{A}_\chi]$, therefore $(\mathbf{A}_\chi)_{\chi^{-1}} = \mathbf{A}$. The main difference is that in $G^2[\mathbf{A}]$ the convolution does not define a subgroup structure for its 2-cocycles.

3.3.2. Gauge transformations. Let us call \mathbb{T}_ψ the twist transformation related to $\psi \in \mathfrak{Z}^2$ over a given quantum space. The results of the previous subsection imply the twist transformations, associated to an inclusion of vector spaces $\mathbf{A}_1 \subset \mathbf{A}$, define a groupoid with (partial) composition $\mathbb{T}_\psi \mathbb{T}_\varphi = \mathbb{T}_{\psi * \varphi}$ and inversion $\mathbb{T}_\psi^{-1} = \mathbb{T}_{i\psi}$. As a generalization of this fact, let us consider the following definition.

Definition 4. We shall say a couple of quantum spaces \mathcal{A} and \mathcal{B} are **twist or gauge related**, namely $\mathcal{A} \sim \mathcal{B}$, if there exists an \mathcal{A} -admissible $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$ and an isomorphism of quantum spaces $\alpha : \mathcal{A}_\psi \simeq \mathcal{B}$. The pairs (α, ψ) will be called **gauge transformations** between \mathcal{A} and \mathcal{B} . ■

Proposition 15. The twist relation is an equivalence relation.

Proof. Reflexivity: Since $\mathbb{I}^{\otimes 2} \in \mathfrak{Z}^2[\mathbf{A}_1]$ is trivially \mathcal{A} -admissible, the identity morphism insures $\mathcal{A} \sim \mathcal{A}$.

Symmetry: Suppose $\mathcal{A} \sim \mathcal{B}$ via a cochain ψ and an isomorphism α . Let θ be the primitive of ψ , and consider the 2-cocycle $i\psi = (\theta \otimes \theta)^{-1} \psi^{-1} (\theta \otimes \theta) \in \mathfrak{Z}^2[\mathbf{A}_1]$. Since the restriction $\alpha_1 = \alpha|_{\mathbf{A}_1}$ defines an isomorphism $\mathbf{A}_1 \simeq \mathbf{B}_1$, through the functor given in §2.3 we have a bijection $\mathfrak{Z}^2[\mathbf{A}_1] \simeq \mathfrak{Z}^2[\mathbf{B}_1]$, such that

$$\psi \mapsto \psi^{\alpha_1} = (\alpha_1^{\otimes 2} \otimes \alpha_1^{\otimes 2}) \psi (\alpha_1^{\otimes 2} \otimes \alpha_1^{\otimes 2})^{-1}.$$

In particular, $i\psi^{\alpha_1} = (\alpha_1^{\otimes 2} \otimes \alpha_1^{\otimes 2}) i\psi (\alpha_1^{\otimes 2} \otimes \alpha_1^{\otimes 2})^{-1} \in \mathfrak{Z}^2[\mathbf{B}_1]$. Using that $\alpha \Pi_{\mathbf{A}} \theta = \Pi_{\mathbf{B}} \alpha_1^{\otimes 2}$ and $\Psi \Pi_{\mathbf{A}}^{\otimes 2} = \Pi_{\mathbf{A}}^{\otimes 2} \psi$, we have

$$\Pi_{\mathbf{B}}^{\otimes 2} i\psi^{\alpha_1} = (\alpha \otimes \alpha) \Psi^{-1} (\alpha \otimes \alpha)^{-1} \Pi_{\mathbf{B}}^{\otimes 2}.$$

Thus, $i\psi^{\alpha_1}$ is \mathcal{B} -admissible and defines the twisted product

$$m_{\mathbf{B}} (\alpha \otimes \alpha) \Psi^{-1} (\alpha \otimes \alpha)^{-1}.$$

On the other hand, since $m_{\mathbf{B}} (\alpha \otimes \alpha) = \alpha m_{\mathbf{A}} \Psi$,

$$\alpha^{-1} \left[m_{\mathbf{B}} (\alpha \otimes \alpha) \Psi^{-1} (\alpha \otimes \alpha)^{-1} \right] = m_{\mathbf{A}} (\alpha \otimes \alpha)^{-1},$$

therefore α^{-1} defines a quantum space isomorphism $\mathcal{B}_{i\psi^{\alpha_1}} \simeq \mathcal{A}$, and consequently $\mathcal{B} \sim \mathcal{A}$.

Transitivity: Consider another quantum space \mathcal{C} such that $\mathcal{B} \sim \mathcal{C}$ via a cochain φ and an isomorphism β . Using again $\alpha \Pi_{\mathbf{A}} \theta = \Pi_{\mathbf{B}} \alpha_1^{\otimes 2}$ and $\Psi \Pi_{\mathbf{A}}^{\otimes 2} = \Pi_{\mathbf{A}}^{\otimes 2} \psi$, it can be seen that

$$\psi \star \varphi^{\alpha_1^{-1}} = \psi (\theta \otimes \theta) (\alpha_1^{\otimes 2} \otimes \alpha_1^{\otimes 2})^{-1} \varphi (\alpha_1^{\otimes 2} \otimes \alpha_1^{\otimes 2}) (\theta \otimes \theta)^{-1}$$

is \mathcal{A} -admissible and has a related automorphism $\Psi (\alpha \otimes \alpha)^{-1} \Phi (\alpha \otimes \alpha)$, if $\Phi \Pi_{\mathbf{B}}^{\otimes 2} = \Pi_{\mathbf{B}}^{\otimes 2} \varphi$. It defines on \mathcal{A} the twisted product $m_{\mathbf{A}} \Psi (\alpha \otimes \alpha)^{-1} \Phi (\alpha \otimes \alpha)$. Then,

$$\begin{aligned} \beta \alpha \left[m_{\mathbf{A}} \Psi (\alpha \otimes \alpha)^{-1} \Phi (\alpha \otimes \alpha) \right] &= \beta m_{\mathbf{B}} \Phi (\alpha \otimes \alpha) \\ &= m_{\mathbf{C}} (\beta \alpha \otimes \beta \alpha), \end{aligned}$$

where we have used $\beta m_{\mathbf{B}} \Phi = m_{\mathbf{C}} (\beta \otimes \beta)$. It follows that $\mathcal{A} \sim \mathcal{C}$. ■

Denoting by $\mathbb{T}_{(\alpha, \psi)}$ the gauge transformation defined by (α, ψ) , the theorem above says those transformations form a groupoid (or a category based on quantum spaces) with composition and inverse

$$\mathbb{T}_{(\beta, \varphi)} \mathbb{T}_{(\alpha, \psi)} \doteq \mathbb{T}_{(\beta, \varphi) \star (\alpha, \psi)}, \quad \mathbb{T}_{(\alpha, \psi)}^{-1} \doteq \mathbb{T}_{i(\alpha, \psi)},$$

being

$$(\beta, \varphi) \star (\alpha, \psi) \doteq (\beta \alpha, \psi \star \varphi^{\alpha_1^{-1}}) \quad \text{and} \quad i(\alpha, \psi) \doteq (\alpha^{-1}, i\psi^{\alpha_1}).$$

As examples of gauge equivalence we have $\underline{\text{hom}}^{\mathbb{T}}[\mathcal{B}, \mathcal{A}] \sim \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$ for every pair \mathcal{B}, \mathcal{A} in CA, and $\mathcal{A} \circ_{\tau} \mathcal{B} \sim \mathcal{A} \circ \mathcal{B}$ for every STTP of \mathcal{A} and \mathcal{B} .

Geometrically, we are defining an equivalence relation among non commutative algebraic varieties in QLS.

A characterization of gauge equivalence between conic quantum spaces can be given in terms of their corresponding ideals. More precisely,

Theorem 12. Let \mathcal{A}, \mathcal{B} be objects of CA, with related ideals \mathbf{I} and \mathbf{J} , respectively. $\mathcal{A} \sim \mathcal{B}$ iff there exists an homogeneous (of degree zero) linear isomorphism $\vartheta : \mathbf{A}_1^{\otimes} \simeq \mathbf{B}_1^{\otimes}$ such that

$$(3.24) \quad \vartheta (\mathbf{A}_1^{\otimes r} \cdot \mathbf{I}_s + \mathbf{I}_r \cdot \mathbf{A}_1^{\otimes s}) = \mathbf{B}_1^{\otimes r} \cdot \mathbf{J}_s + \mathbf{J}_r \cdot \mathbf{B}_1^{\otimes s}; \quad r, s \in \mathbb{N}_0.$$

Proof. Suppose \mathcal{A} is gauge equivalent to \mathcal{B} , i.e. there exists an \mathcal{A} -admissible $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$ and an isomorphism of quantum spaces $\alpha : \mathcal{A}_\psi \simeq \mathcal{B}$. If $\psi = \partial\theta$, from **Lemma 3**, admissibility condition is equivalent to

$$(3.25) \quad \theta^{-1}(\mathbf{A}_1^{\otimes r} \cdot \mathbf{I}_s + \mathbf{I}_r \cdot \mathbf{A}_1^{\otimes s}) = \mathbf{A}_1^{\otimes r} \cdot \theta^{-1}(\mathbf{I}_s) + \theta^{-1}(\mathbf{I}_r) \cdot \mathbf{A}_1^{\otimes s}.$$

Because α defines an algebra homomorphism $\mathbf{A}_\psi \simeq \mathbf{B}$, that is to say

$$\alpha_1^{\otimes}(\mathbf{I}_{\psi,r}) = \alpha_1^{\otimes} \theta^{-1}(\mathbf{I}_r) = \mathbf{J}_r,$$

we have

$$(3.26) \quad \begin{aligned} & \alpha_1^{\otimes} \theta^{-1}(\mathbf{A}_1^{\otimes r} \cdot \mathbf{I}_s + \mathbf{I}_r \cdot \mathbf{A}_1^{\otimes s}) = \alpha_1^{\otimes}(\mathbf{A}_1^{\otimes r} \cdot \theta^{-1}(\mathbf{I}_s) + \theta^{-1}(\mathbf{I}_r) \cdot \mathbf{A}_1^{\otimes s}) \\ & = \alpha_1^{\otimes}(\mathbf{A}_1^{\otimes r}) \cdot \alpha_1^{\otimes} \theta^{-1}(\mathbf{I}_s) + \alpha_1^{\otimes} \theta^{-1}(\mathbf{I}_r) \cdot \alpha_1^{\otimes}(\mathbf{A}_1^{\otimes s}) \\ & = \mathbf{B}_1^{\otimes r} \cdot \mathbf{J}_s + \mathbf{J}_r \cdot \mathbf{B}_1^{\otimes s}, \end{aligned}$$

The linear map $\vartheta = \alpha_1^{\otimes} \theta^{-1} : \mathbf{A}_1^{\otimes} \rightarrow \mathbf{B}_1^{\otimes}$ is obviously the map we are looking for.

Reciprocally, consider an homogeneous linear bijection $\vartheta : \mathbf{A}_1^{\otimes} \simeq \mathbf{B}_1^{\otimes}$ satisfying (3.24). Given a basis $\{a_i\}$ of \mathbf{A}_1 , if $\vartheta(a_i) = b_i$, then $\{b_i\}$ is a basis of \mathbf{B}_1 , and we can write

$$\vartheta(a_{i_1} \dots a_{i_n}) = (\vartheta_n)_{i_1 \dots i_n}^{j_1 \dots j_n} b_{j_1} \dots b_{j_n}; \quad \vartheta(1) = 1.$$

The assignment $\alpha_1 = \vartheta|_{\mathbf{A}_1} : a_i \mapsto b_i$ defines a linear bijection, and ϑ can be decomposed as $\vartheta = \alpha_1^{\otimes} \theta^{-1}$ with

$$\theta^{-1}(a_{i_1} \dots a_{i_n}) = (\vartheta_n)_{i_1 \dots i_n}^{j_1 \dots j_n} a_{j_1} \dots a_{j_n}; \quad \theta(1) = 1.$$

In particular, since $(\vartheta_1)_i^j = \delta_i^j$ and $\theta(1) = 1$, then $\theta_{0,1} = \mathbb{I}_{0,1}$ and, as a consequence, $\theta \in \mathfrak{P}^1[\mathbf{A}_1]$. Let us consider the counital 2-cocycle $\psi = \partial\theta$. We shall see ψ is \mathcal{A} -admissible and α_1^{\otimes} defines an isomorphism $\mathcal{A}_\psi \simeq \mathcal{B}$.

Multiplying Eq. (3.24) by $(\alpha_1^{\otimes})^{-1}$ to the left, we obtain precisely Eq. (3.25). But this is the admissibility condition for ψ , provided $\psi_{r,s} \approx \theta_{r+s}(\theta_r \otimes \theta_s)^{-1}$. Now, using again (3.24), but in the form (3.26) and fixing $s = 0$, equation $\alpha_1^{\otimes}(\mathbf{I}_{\psi,r}) = \mathbf{J}_r$ follows. ■

In the general case we can say,

Proposition 16. *Given $\mathcal{A}, \mathcal{B} \in \text{FGA}$, with related ideals \mathbf{I} and \mathbf{J} , if $\mathcal{A} \sim \mathcal{B}$, then there exists an homogeneous (of degree zero) linear bijection $\vartheta : \mathbf{A}_1^{\otimes} \simeq \mathbf{B}_1^{\otimes}$ such that $\vartheta(\mathbf{I}) = \mathbf{J}$.*

Proof. If $\mathcal{A} \sim \mathcal{B}$ through $\psi = \partial\theta \in \mathfrak{Z}^2[\mathbf{A}_1]$ and an isomorphism $\alpha : \mathcal{A}_\psi \simeq \mathcal{B}$, then $\vartheta = \alpha_1^{\otimes} \theta^{-1}$ is an homogeneous linear isomorphism such that $\vartheta(\mathbf{I}) = \mathbf{J}$. ■

Thus, gauge equivalent non commutative spaces in QLS have isomorphic defining ideals.

4. TWISTING AND FUNCTORS ON CA

In this chapter we mainly deal with conic quantum spaces, analyzing the relationship between certain functors defined on their category, and quasicomplexes naturally related to them. For instance, $\mathfrak{C}^\bullet[\mathbf{A}_1]$ and $\mathfrak{C}^\bullet[\mathbf{A}_1^*]$ are quasicomplexes naturally related to the functor $! : \mathcal{A} \mapsto \mathcal{A}^!$, while $\mathfrak{C}^\bullet[\mathbf{A}_1]$, $\mathfrak{C}^\bullet[\mathbf{B}_1]$ and $\mathfrak{C}^\bullet[\mathbf{A}_1 \otimes \mathbf{B}_1]$ are related to bifunctors \circ , \bullet and \odot . We shall study the following problems. Suppose a twisting \mathcal{A}_φ of \mathcal{A} is given (of course, $\varphi \in \mathfrak{Z}^2[\mathbf{A}_1]$): when can we insure there exists $\chi \in \mathfrak{Z}^2[\mathbf{A}_1^*]$ such that $(\mathcal{A}_\varphi)^! = (\mathcal{A}^!)_\chi$? On the other hand, given another twisted quantum space, namely \mathcal{B}_ϕ (now $\phi \in \mathfrak{Z}^2[\mathbf{B}_1]$): can we write

$$\mathcal{A}_\varphi \circ \mathcal{B}_\phi = (\mathcal{A} \circ \mathcal{B})_\kappa, \quad \circ = \circ, \bullet \text{ or } \odot,$$

for some $\kappa \in \mathfrak{Z}^2[\mathbf{A}_1 \otimes \mathbf{B}_1]$?

In the following, we answer these questions and other corresponding to the functors \triangleleft , \triangleright and \diamond . In relation to latter functors, we study the internal coHom objects of twisted quantum spaces, comparing them with the untwisted ones.

4.1. 2nd admissibility condition. For conic quantum spaces, a stronger condition than admissibility is defined. Its consequences are analyzed along all subsequent subsections.

Definition 5. *Let \mathcal{A} be a conic quantum space with related ideal $\mathbf{I} = \bigoplus_{n \geq 2} \mathbf{I}_n$. We shall call **2nd \mathcal{A} -admissible**, or **2nd admissible** for \mathcal{A} , the n -cochains satisfying*

$$(4.1) \quad \begin{aligned} & \psi(\mathbf{A}_1^{\otimes p_1} \otimes \dots \otimes (\mathbf{A}_1^{\otimes p_i} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) \otimes \dots \otimes \mathbf{A}_1^{\otimes p_n}) \\ & = \mathbf{A}_1^{\otimes p_1} \otimes \dots \otimes (\mathbf{A}_1^{\otimes p_i} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) \otimes \dots \otimes \mathbf{A}_1^{\otimes p_n}, \end{aligned}$$

for all $p_k, q, r \in \mathbb{N}_0$, $k, i = 1 \dots n$. ■

For 2-cochains, 2nd admissibility is equivalent to the inclusions

$$(4.2) \quad \begin{aligned} \psi (\mathbf{A}_1^{\otimes s} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r})) &\subset \mathbf{A}_1^{\otimes s} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}), \\ \psi ((\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) \otimes \mathbf{A}_1^{\otimes s}) &\subset (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) \otimes \mathbf{A}_1^{\otimes s}; \end{aligned}$$

and **Lemma 3** translates into the next result.

Lemma 4. *Under the conditions of above definition, $\psi = \partial\theta$ is 2nd admissible iff*

$$(4.3) \quad \theta^{-1} (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes p} \cdot \theta^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}.$$

Proof. If (4.3) holds for the primitive θ of ψ , we have, putting $p = x + y$,

$$\theta^{-1} (\mathbf{A}_1^{\otimes x} \cdot \mathbf{A}_1^{\otimes y} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes x} \cdot \mathbf{A}_1^{\otimes y} \cdot \theta^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}.$$

But $\mathbf{A}_1^{\otimes y} \cdot \theta^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r} = \theta^{-1} (\mathbf{A}_1^{\otimes y} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r})$ and $\mathbf{A}_1^{\otimes x} = \theta^{-1} (\mathbf{A}_1^{\otimes x})$, thus

$$\theta^{-1} (\mathbf{A}_1^{\otimes x} \cdot (\mathbf{A}_1^{\otimes y} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r})) = \theta^{-1} (\mathbf{A}_1^{\otimes x}) \cdot \theta^{-1} (\mathbf{A}_1^{\otimes y} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}),$$

and using the fact that $\psi = \partial\theta$, the first part of (4.2) follows. In a similar way the second part can be shown, and 2nd admissibility is fulfilled. The converse follows in a similar way. ■

Clearly, if ψ is 2nd admissible, then is admissible. Furthermore, the 2nd admissible cochains form a subgroup of the admissible ones.

4.1.1. *Restricted gauge equivalence.* For 2-cocycles we have, in relation to the product \star in \mathfrak{Z}^2 , that:

Proposition 17. *Let ψ and φ be a couple of counital 2-cocycles, 2nd admissible for \mathcal{A} and \mathcal{A}_ψ , respectively. Then $\psi \star \varphi$ is 2nd \mathcal{A} -admissible. Furthermore $i\psi$, the inverse of ψ under the product \star , is 2nd admissible for \mathcal{A}_ψ .*

Proof. Suppose $\psi = \partial\lambda$ and $\varphi = \partial\chi$. From lemma above, λ and χ satisfies

$$(4.4) \quad \lambda^{-1} (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes p} \cdot \lambda^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}$$

and

$$\chi^{-1} (\mathbf{A}_1^{\otimes p} \cdot \lambda^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes p} \cdot (\lambda\chi)^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r},$$

since $\lambda^{-1} (\mathbf{I})$ is the ideal related to \mathcal{A}_ψ . In consequence,

$$(\lambda\chi)^{-1} (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes p} \cdot (\lambda\chi)^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}.$$

On the other hand, $\psi \star \varphi = \partial\lambda \star \partial\chi = \partial(\lambda\chi)$, hence $\psi \star \varphi$ has a primitive satisfying (4.3). Therefore, $\psi \star \varphi$ is 2nd admissible for \mathcal{A} .

For the second claim of our proposition, using that $i\psi = \partial(\lambda^{-1})$ and (from Eq. (4.4))

$$\begin{aligned} \lambda (\mathbf{A}_1^{\otimes p} \cdot \lambda^{-1} (\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}) &= \lambda (\lambda^{-1} (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r})) \\ &= \mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r} = \mathbf{A}_1^{\otimes p} \cdot \lambda (\lambda^{-1} (\mathbf{I}_q)) \cdot \mathbf{A}_1^{\otimes r}, \end{aligned}$$

the 2nd \mathcal{A}_ψ -admissibility of $i\psi$ is immediate. ■

Therefore, in terms of 2nd admissible cocycles a restricted twist equivalence can be defined. A characterization of equivalent quantum spaces through this gauge transformations follow from the proof of **Theor. 12**, and says:

Theorem 13. *Let \mathcal{A}, \mathcal{B} be objects of CA, with related ideals \mathbf{I} and \mathbf{J} , respectively. $\mathcal{A} \sim \mathcal{B}$ iff there exists an homogeneous (of degree zero) linear isomorphism $\vartheta : \mathbf{A}_1^\otimes \simeq \mathbf{B}_1^\otimes$ such that*

$$\vartheta (\mathbf{A}_1^{\otimes r} \cdot \mathbf{I}_s \cdot \mathbf{A}_1^{\otimes t}) = \mathbf{B}_1^{\otimes r} \cdot \mathbf{J}_s \cdot \mathbf{B}_1^{\otimes t}; \quad r, s, t \in \mathbb{N}_0. \quad \blacksquare$$

By last, consider the following property of 2nd admissible 2-cocycles.

Proposition 18. *If a pair of 2nd admissible 2-cocycles are cohomologous through an admissible 1-cochain θ , then θ is 2nd admissible.*

Proof. Let $\psi = \partial\lambda$ and $\varphi = \partial\chi$ be cohomologous 2-cocycles through an admissible cochain θ . We know from **Prop. 11** that $\theta = \lambda\omega\chi^{-1}$, with ω a 1-cocycle. Using **Lemma 4** for χ and then the admissibility of θ (and the fact that ω is a 1-cocycle),

$$\begin{aligned} \lambda\omega\chi^{-1}(\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) &= \lambda(\mathbf{A}_1^{\otimes p} \cdot \omega\chi^{-1}(\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}) \\ &= \lambda(\mathbf{A}_1^{\otimes p} \cdot \lambda^{-1}(\lambda\omega\chi^{-1})(\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}) = \lambda(\mathbf{A}_1^{\otimes p} \cdot \lambda^{-1}(\mathbf{I}_q) \cdot \mathbf{A}_1^{\otimes r}). \end{aligned}$$

Using same lemma for λ we have $\theta(\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}$, as we have claimed. ■

From that, **Theor. 9** can be rephrased in terms of 2nd admissibility too.

4.1.2. *The (anti)bicharacter case.* Examples of 2nd admissible cochains are the admissible (anti)bicharacters.

Proposition 19. *Let $\psi \in \mathfrak{C}^2[\mathbf{A}_1]$ be an (anti)bicharacter. ψ is admissible iff is 2nd admissible.* ■

That follows from proposition below.

Proposition 20. *Let $\psi \in \mathfrak{C}^2[\mathbf{A}_1]$ be an (anti)bicharacter for which there exists a graded vector space*

$$\mathbf{S} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{S}_n \subset \mathbf{A}_1^{\otimes} = \bigoplus_{n \in \mathbb{N}_0} \mathbf{A}_1^{\otimes n}; \quad \mathbf{S}_{0,1} \doteq \{0\},$$

such that

$$(4.5) \quad \psi(\mathbf{A}_1^{\otimes m} \otimes \mathbf{S}_n + \mathbf{S}_m \otimes \mathbf{A}_1^{\otimes n}) = \mathbf{A}_1^{\otimes m} \otimes \mathbf{S}_n + \mathbf{S}_m \otimes \mathbf{A}_1^{\otimes n}.$$

Then, for all $p, q, r, u \in \mathbb{N}_0$, we have

$$(4.6) \quad \begin{aligned} \psi((\mathbf{A}_1^{\otimes p} \cdot \mathbf{S}_q \cdot \mathbf{A}_1^{\otimes r}) \otimes \mathbf{A}_1^{\otimes u}) &= (\mathbf{A}_1^{\otimes p} \cdot \mathbf{S}_q \cdot \mathbf{A}_1^{\otimes r}) \otimes \mathbf{A}_1^{\otimes u}, \\ \psi(\mathbf{A}_1^{\otimes u} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{S}_q \cdot \mathbf{A}_1^{\otimes r})) &= \mathbf{A}_1^{\otimes u} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{S}_q \cdot \mathbf{A}_1^{\otimes r}); \end{aligned}$$

and if ψ is a 2-cocycle with primitive θ ,

$$(4.7) \quad \theta^{-1}(\mathbf{A}_1^{\otimes p} \cdot \mathbf{S}_q \cdot \mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes p} \cdot \theta^{-1}(\mathbf{S}_q) \cdot \mathbf{A}_1^{\otimes r}.$$

Proof. Consider an element $a \in \mathbf{A}_1$ and $s \in \mathbf{S}_n$. Since Eq. (4.5),

$$\psi(a \otimes s) = \psi_{1,n}(a \otimes s) \in \mathbf{A}_1 \otimes \mathbf{S}_n + \mathbf{S}_1 \otimes \mathbf{A}_1^{\otimes n}.$$

But $\mathbf{S}_1 = \{0\}$, hence

$$(4.8) \quad \psi_{1,n}(\mathbf{A}_1 \otimes \mathbf{S}_n) = \mathbf{A}_1 \otimes \mathbf{S}_n \quad \text{or} \quad \psi_{1,n}(\mathbf{A}_1 \otimes \mathbf{S}_n) = \mathbf{A}_1 \otimes \mathbf{S}_n.$$

Now, consider an element $a \otimes b = a \cdot b \in \mathbf{A}_1^{\otimes 2}$. Suppose ψ is a bicharacter. From the first part of Eq. (2.25) we have

$$\begin{aligned} \psi_{2,n}((a \cdot b) \otimes s) &= \psi_{1+1,n}((a \cdot b) \otimes s) \\ &= (\mathbb{I} \otimes \psi_{1,n})(\mathbb{I} \otimes f_{1,n}^{-1})(\psi_{1,n}(a \otimes s) \otimes b), \end{aligned}$$

and from Eq. (4.8) it follows that $\psi(\mathbf{A}_1^{\otimes 2} \otimes \mathbf{S}_n) = \mathbf{A}_1^{\otimes 2} \otimes \mathbf{S}_n$. An inductive reasoning shows that

$$\psi(\mathbf{A}_1^{\otimes m} \otimes \mathbf{S}_n) = \psi_{m,n}(\mathbf{A}_1^{\otimes m} \otimes \mathbf{S}_n) = \mathbf{A}_1^{\otimes m} \otimes \mathbf{S}_n$$

for all n, m . Now, consider an element $c \otimes (a \cdot s) \in \mathbf{A}_1^{\otimes m} \otimes \mathbf{A}_1 \cdot \mathbf{S}_n$. Using the second part of Eq. (2.25),

$$\psi_{m,1+n}(c \otimes (a \cdot s)) = (\psi_{m,1} \otimes \mathbb{I}_n)(\mathbb{I}_m \otimes f_{1,n}^{-1})(\psi_{m,n}(c \otimes s) \otimes a),$$

hence $\psi(\mathbf{A}_1^{\otimes m} \otimes \mathbf{A}_1 \cdot \mathbf{S}_n) = \mathbf{A}_1^{\otimes m} \otimes \mathbf{A}_1 \cdot \mathbf{S}_n$. Analogous arguments, followed by inductive reasoning, lead us finally to the equalities (4.6).

To prove the second part of the lemma, i.e. Eq. (4.7), recall that a 2-cocycle ψ and its primitive θ are related by the equation

$$\theta_{m+n}^{-1} \approx (\theta_m^{-1} \otimes \theta_n^{-1}) \psi_{m,n}^{-1}.$$

Thus, the previous result implies $\theta^{-1}(\mathbf{S}_m \cdot \mathbf{A}_1^{\otimes n}) = \theta^{-1}(\mathbf{S}_m) \cdot \theta^{-1}(\mathbf{A}_1^{\otimes n})$, and the other equalities follow in a similar way. This ends our proof. ■

Each bicharacter $\varsigma_{\mathcal{A}}$ appearing in twisted coHom objects (see (3.18)) is admissible, *ipso facto* 2nd admissible, thanks to $\sigma_{\mathcal{A}}$ defines an automorphism. In fact,

$$\begin{aligned} \varsigma_{\mathcal{A}}(\mathbf{A}_1^{\otimes s} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r})) &= \mathbf{A}_1^{\otimes s} \otimes [\sigma_{\mathcal{A}}^{-s}]^{\otimes} (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}) \\ &= \mathbf{A}_1^{\otimes s} \otimes [\sigma_{\mathcal{A}}^{-s}]^{\otimes} (\mathbf{A}_1^{\otimes p}) \cdot [\sigma_{\mathcal{A}}^{-s}]^{\otimes} (\mathbf{I}_q) \cdot [\sigma_{\mathcal{A}}^{-s}]^{\otimes} (\mathbf{A}_1^{\otimes r}) = \mathbf{A}_1^{\otimes s} \otimes \mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}. \end{aligned}$$

4.1.3. *Quasicomplexes of 2nd admissible cochains.* From Eq. (4.1), easy calculations show that the coface operators and codegeneracies on \mathbf{C}^\bullet send the subgroup of 2nd admissible cochains to itself. Thus, we can define for each conic quantum space $\mathcal{A} = (\mathbf{A}_1, \mathbf{A})$ a subquasicomplex $\mathbf{C}^\bullet[\mathcal{A}]$ of $\mathbf{C}^\bullet[\mathbf{A}_1]$ formed out by 2nd \mathcal{A} -admissible cochains. We are not going to employ explicitly this quasicomplex in any calculation. However, for the sake of completeness, we enumerate some of its properties bellow (without proof).

- As in §2.3, the cosimplicial objects $[n+1] \mapsto \mathbf{C}^n[\mathcal{A}]$ and $[n+1] \mapsto \mathbf{C}^n[\mathcal{B}]$ are naturally equivalent when $\mathcal{A} \simeq \mathcal{B}$.

- The map $\mathcal{A} \mapsto \mathbf{C}^\bullet[\mathcal{A}]$ defines a functor $\mathbf{C}^\bullet : \mathcal{G}[\mathbf{CA}] \rightarrow \text{Grp}_{*q}$ and a morphism between the groupoids $\mathcal{G}[\mathbf{CA}]$ and $\mathcal{G}[\text{Grp}_{*q}]$ such that to every isomorphism $\alpha : \mathcal{A} \simeq \mathcal{B}$ it assigns the isomorphism of quasicomplexes α^\bullet defined as in Eq. (2.28), replacing f by α_1 .

- The subgroups of counital cochains give rise to a subquasicomplex $\mathfrak{C}^\bullet[\mathcal{A}]$ satisfying Eq. (2.22).

- The first and second cohomology spaces, namely $H^{1,2}[\mathcal{A}]$, can be defined as in §3.2. Now, they are usual cohomological spaces in the sense that are given by quotients $Z^{1,2}[\mathcal{A}] / \sim_{\text{Coh}}$. It follows that

$$H^1[\mathcal{A}] = Z^1[\mathcal{A}] \simeq \text{Aut}_{\text{CA}}[\mathcal{A}] \simeq H_{\mathcal{A}}^1[\mathbf{A}_1],$$

and from **Prop. 18** the inclusion $H^2[\mathcal{A}] \subset H_{\mathcal{A}}^2[\mathbf{A}_1]$ is immediate.

- From Eq. (2.44) (putting $\mathbf{V} = \mathbf{A}_1$), it follows by restriction the isomorphism

$$(4.9) \quad \mathbf{C}^\bullet[\mathcal{A}] \simeq \mathbf{G}^\bullet[\underline{\text{end}}[\mathcal{A}]] = \mathbf{G}^\bullet[\mathcal{A} \triangleright \mathcal{A}].$$

In particular, suppose \mathcal{A} is quadratic and consider the algebras $\underline{e}[\mathcal{A}]$ and $A(R)$ defined in [12] and [14],¹³ respectively. The well known associated epimorphisms $A(R) \leftarrow \underline{\text{end}}[\mathcal{A}] \rightarrow \underline{e}[\mathcal{A}]$, which follows from initiality of $\underline{\text{end}}[\mathcal{A}]$, gives rise to monics

$$\mathbf{G}^\bullet[A(R)] \hookrightarrow \mathbf{G}^\bullet[\underline{\text{end}}[\mathcal{A}]] \hookrightarrow \mathbf{G}^\bullet[\underline{e}[\mathcal{A}]].$$

Thus, the bialgebra twisting of $A(R)$ and $\underline{e}[\mathcal{A}]$ are given by a particular subgroup of 2nd admissible twisting of the quantum space \mathcal{A} .

Another properties of $\mathbf{C}^\bullet[\mathcal{A}]$, in relation with the involution and the product between quasicomplexes in $\mathbf{C}^\bullet[\text{Vct}]$, will be briefly commented in the following section.

4.2. Coadjoints, products and internal coHom objects.

4.2.1. *Dual quantum spaces and coadjoint cochains.* Let us first enunciate the following well-known result (without proof).

Lemma 5. *Consider a pair of finite dimensional \mathbb{k} -vector spaces \mathbf{U} and \mathbf{V} , supplied with a non degenerated pairing $\mathbf{U} \times \mathbf{V} \rightarrow \mathbb{k}$. Let α be an automorphism in \mathbf{V} . Then, for every subset $\mathbf{S} \subset \mathbf{V}$ we have the equality of vector spaces¹⁴*

$$\alpha(\mathbf{S})^\perp = \alpha^{*-1}(\mathbf{S}^\perp),$$

where α^* denotes the (transpose) automorphism induced in \mathbf{U} by the mentioned pairing. ■

Consider a conic quantum space \mathcal{A} and a counital \mathcal{A} -admissible 2-cocycle $\psi = \partial\theta \in \mathfrak{Z}^2[\mathbf{A}_1]$. The ideal \mathbf{I}_ψ related to \mathcal{A}_ψ is determined by $\mathbf{I}_{\psi,n} = \theta^{-1}(\mathbf{I}_n)$, hence (from lemma above)

$$(\mathbf{I}_{\psi,n})^\perp = \theta^{-1}(\mathbf{I}_n)^\perp = \theta^*(\mathbf{I}_n^\perp),$$

and recalling Eq. (1.4), $(\mathbf{I}_\psi)^\dagger = I \left[\bigoplus_{n \geq 2} (\mathbf{I}_{\psi,n})^\perp \right] = I \left[\bigoplus_{n \geq 2} \theta^*(\mathbf{I}_n^\perp) \right]$. On the other hand, consider the 2-cocycle¹⁵

$$\psi^{*-1} = \psi^\dagger = (\partial\theta)^\dagger = \partial^\dagger(\theta^\dagger) \in \mathfrak{Z}^2[\mathbf{A}_1^*] = \mathfrak{Z}^2[\mathbf{A}_1]^\dagger.$$

If ψ^\dagger were \mathcal{A}^\dagger -admissible, it follows from **Prop. 12** (for $\mathbf{S} = \bigoplus_{n \geq 2} \mathbf{I}_n^\perp$) that the ideal related to $(\mathcal{A}^\dagger)_{\psi^\dagger}$ would be

$$(\mathbf{I}^\dagger)_{\psi^\dagger} = \theta^*(\mathbf{I}^\dagger) = I \left[\bigoplus_{n \geq 2} \theta^*(\mathbf{I}_n^\perp) \right].$$

That is to say, $(\mathcal{A}_\psi)^\dagger$ and $(\mathcal{A}^\dagger)_{\psi^\dagger}$ coincide. The problem is to insure the admissibility of ψ^\dagger . The things change when we suppose ψ is 2nd admissible or, in particular, if ψ is an admissible (anti)bicharacter.

Proposition 21. *If $\psi \in \mathfrak{C}^n[\mathbf{A}_1]$ is 2nd \mathcal{A} -admissible, then $\psi^\dagger \in \mathfrak{C}^n[\mathbf{A}_1]^\dagger$ is 2nd \mathcal{A}^\dagger -admissible. For $\mathcal{A} \in \text{CA}^m$, the converse is also valid.*

¹³If the ideal related to \mathcal{A} is generated by elements $R_{ij}^{kl} a_k a_l$, being R_{ij}^{kl} the coefficients of a map $R : \mathbf{A}_1^{\otimes 2} \rightarrow \mathbf{A}_1^{\otimes 2}$, then the ideal related to $\text{frit}[\mathcal{A}, R]$ is generated by $R_{ij}^{kl} z_k^i z_l^m - z_i^k z_j^l R_{kl}^{nm}$.

¹⁴As usual, given a subset $\mathbf{S} \subset \mathbf{V}$, \mathbf{S}^\perp is the vector space $\mathbf{S}^\perp = \{x \in \mathbf{U} : \langle x, y \rangle = 0, \forall y \in \mathbf{S}\}$.

¹⁵Recall the isomorphism of quasicomplexes $\mathbf{C}^\bullet[\mathbf{V}] \simeq \mathbf{C}^\bullet[\mathbf{V}]^\dagger = \mathbf{C}^\bullet[\mathbf{V}^*]$.

Proof. We just consider $n = 2$, since the other cases can be shown in a similar way. From **Lemma 5**,

$$\begin{aligned} \psi^! (\mathbf{A}_1^{*\otimes u} \otimes (\mathbf{A}_1^{*\otimes p} \cdot \mathbf{I}_q^\perp \cdot \mathbf{A}_1^{*\otimes r})) &= \psi^! \left((\mathbf{A}_1^{\otimes u} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}))^\perp \right) \\ &= \psi (\mathbf{A}_1^{\otimes u} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}))^\perp, \end{aligned}$$

and from 2nd \mathcal{A} -admissibility of ψ , i.e.

$$(4.10) \quad \psi (\mathbf{A}_1^{\otimes u} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r})) = \mathbf{A}_1^{\otimes u} \otimes (\mathbf{A}_1^{\otimes p} \cdot \mathbf{I}_q \cdot \mathbf{A}_1^{\otimes r}),$$

we have

$$\psi^! (\mathbf{A}_1^{*\otimes u} \otimes (\mathbf{A}_1^{*\otimes p} \cdot \mathbf{I}_q^\perp \cdot \mathbf{A}_1^{*\otimes r})) = \mathbf{A}_1^{*\otimes u} \otimes (\mathbf{A}_1^{*\otimes p} \cdot \mathbf{I}_q^\perp \cdot \mathbf{A}_1^{*\otimes r}).$$

Straightforwardly, because $\mathbf{I}_n^\perp = \sum_{r=2}^n \sum_{i=0}^{n-r} \mathbf{A}_1^{*\otimes n-r-i} \cdot \mathbf{I}_r^\perp \cdot \mathbf{A}_1^{*\otimes i}$, $\psi^!$ is 2nd $\mathcal{A}^!$ -admissible. Now, let us shown the other claim. Suppose $\mathcal{A} \in \text{CA}^m$. From the last result, if $\psi^! \in \mathfrak{C}^n[\mathbf{A}_1]^!$ is 2nd $\mathcal{A}^!$ -admissible, then $\psi^{\#}$ is 2nd $\mathcal{A}^{\#}$ -admissible. But $\psi^{\#} = \psi$ and the ideal related to $\mathcal{A}^{\#}$ is the same as the one related to \mathcal{A} . So, the proposition have been proven. \blacksquare

In terms of complexes $\mathbf{C}^\bullet[\mathcal{A}]$, last proposition says isomorphism $\mathbf{C}^\bullet[\mathbf{A}_1] \simeq \mathbf{C}^\bullet[\mathbf{A}_1]^!$ gives rise to a monic $\mathbf{C}^\bullet[\mathcal{A}] \hookrightarrow \mathbf{C}^\bullet[\mathcal{A}^!]$ which define a natural transformation $\mathbf{C}^\bullet \rightarrow \mathbf{C}^\bullet!$. In addition, when \mathbf{C}^\bullet is restricted to CA^m , the isomorphisms $\mathbf{C}^\bullet[\mathcal{A}] \simeq \mathbf{C}^\bullet[\mathcal{A}^!]$ and the related natural equivalence $\mathbf{C}^\bullet \simeq \mathbf{C}^\bullet!$ follow.

In resume, we have shown:

Theorem 14. *Let \mathcal{A} be an object in CA . For every counital 2nd \mathcal{A} -admissible $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$, $\psi^!$ is a 2nd $\mathcal{A}^!$ -admissible element of $\mathfrak{Z}^2[\mathbf{A}_1]^!$ and $(\mathcal{A}_\psi)^! = (\mathcal{A}^!)_{\psi^!}$. \blacksquare*

4.2.2. *Compositions of quantum spaces and quasicomplexes.* Given $\mathcal{A}, \mathcal{B} \in \text{FGA}$, consider a couple of related admissible cochains $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$ and $\varphi \in \mathfrak{Z}^2[\mathbf{B}_1]$. From the discussion we made in §2.3.2, they give rise to a 2-cocycle (ψ, φ) in the quasicomplex $\mathfrak{C}^\bullet[\mathbf{A}_1] \times \mathfrak{C}^\bullet[\mathbf{B}_1]$ which, through the monic $j : \mathfrak{C}^\bullet[\mathbf{A}_1] \times \mathfrak{C}^\bullet[\mathbf{B}_1] \hookrightarrow \mathfrak{C}^\bullet[\mathbf{A}_1 \otimes \mathbf{B}_1]$, can be identified with a map (recall Eq. (2.38))

$$j(\psi, \varphi) = \bigoplus_{r,s \in \mathbb{N}_0} \psi_{r,s} \otimes \varphi_{r,s} \in \mathfrak{Z}^2[\mathbf{A}_1 \otimes \mathbf{B}_1].$$

Also, they can be seen as a restriction of $\psi \otimes \varphi$. In these terms, admissibility condition for ψ and φ lead us immediately to $\mathcal{A} \circ \mathcal{B}$ -admissibility of $j(\psi, \varphi) \subset \psi \otimes \varphi$. If $\psi = \partial\theta$ and $\varphi = \partial\chi$, then the primitive of $j(\psi, \varphi)$ is $j(\theta, \chi) = \bigoplus_{r \in \mathbb{N}_0} \theta_r \otimes \chi_r$, and consequently $j(\theta, \chi) \subset \theta \otimes \chi$. From that, it follows the ideal related to $(\mathcal{A} \circ \mathcal{B})_{j(\psi, \varphi)}$ is precisely the one related to $\mathcal{A}_\psi \circ \mathcal{B}_\varphi$. So, we have

Theorem 15. *Given $\mathcal{A}, \mathcal{B} \in \text{FGA}$ and a couple of counital admissible cochains $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$ and $\varphi \in \mathfrak{Z}^2[\mathbf{B}_1]$, then $j(\psi, \varphi) \in \mathfrak{Z}^2[\mathbf{A}_1 \otimes \mathbf{B}_1]$ is $\mathcal{A} \circ \mathcal{B}$ -admissible and $\mathcal{A}_\psi \circ \mathcal{B}_\varphi \simeq (\mathcal{A} \circ \mathcal{B})_{j(\psi, \varphi)}$. \blacksquare*

For other composition of quantum spaces, it is not enough for ψ and φ to be admissible. For instance, the ideal related to $\mathcal{A} \odot \mathcal{B}$, $\bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n \otimes \mathbf{J}_n$, is not preserved by $j(\psi, \varphi)$. Nevertheless, if ψ and φ are 2nd admissible we can show, using similar technics to the ones developed in the previous subsection, that:

- $j(\psi, \varphi)$ is 2nd $\mathcal{A} \odot \mathcal{B}$ -admissible and the related ideal of $(\mathcal{A} \odot \mathcal{B})_{j(\psi, \varphi)}$ is

$$(4.11) \quad j(\theta, \chi)^{-1} \left(\bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n \otimes \mathbf{J}_n \right) = \bigoplus_{n \in \mathbb{N}_0} \theta^{-1}(\mathbf{I}_n) \otimes \chi^{-1}(\mathbf{J}_n);$$

- $j(\psi^!, \varphi)$ is 2nd $\mathcal{A} \triangleright \mathcal{B}$ -admissible and the related ideal of $(\mathcal{A} \triangleright \mathcal{B})_{j(\psi^!, \varphi)}$ is

$$(4.12) \quad \begin{aligned} j(\theta^!, \chi)^{-1} \left(I \left[\bigoplus_{n \in \mathbb{N}_0} \mathbf{I}_n^\perp \otimes \mathbf{J}_n \right] \right) &= I \left[\bigoplus_{n \in \mathbb{N}_0} \theta^{*\perp}(\mathbf{I}_n^\perp) \otimes \chi^{-1}(\mathbf{J}_n) \right] \\ &= I \left[\bigoplus_{n \in \mathbb{N}_0} \theta^{-1}(\mathbf{I}_n)^\perp \otimes \chi^{-1}(\mathbf{J}_n) \right]; \end{aligned}$$

and analogous results for the functors \triangleleft and \diamond . The same can be done if ψ and φ are admissible bicharacters, since, for instance, cochains like $\psi^!$ and $j(\psi, \varphi)$ will also have that property. Note that (4.11) and (4.12) are precisely the ideals related to $\mathcal{A}_\psi \odot \mathcal{B}_\varphi$ and $\mathcal{A}_\psi \triangleright \mathcal{B}_\varphi$, respectively. Summing up,

Theorem 16. *Let $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$ and $\varphi \in \mathfrak{Z}^2[\mathbf{B}_1]$ be 2nd \mathcal{A} and \mathcal{B} -admissible, respectively. Then,*

$$\mathcal{A}_\psi \odot \mathcal{B}_\varphi = (\mathcal{A} \odot \mathcal{B})_{j(\psi, \varphi)}; \quad \mathcal{A}_\psi \triangleright \mathcal{B}_\varphi = (\mathcal{A} \triangleright \mathcal{B})_{j(\psi^!, \varphi)};$$

$$\mathcal{A}_\psi \triangleleft \mathcal{B}_\varphi = (\mathcal{A} \triangleleft \mathcal{B})_{j(\psi, \varphi^!)}; \quad \mathcal{A}_\psi \diamond \mathcal{B}_\varphi = (\mathcal{A} \diamond \mathcal{B})_{j(\psi, \varphi^!)}.$$

And for quadratic and m -th quantum spaces, $\mathcal{A}_\psi \bullet \mathcal{B}_\varphi = (\mathcal{A} \bullet \mathcal{B})_{j(\psi, \varphi)}$. \blacksquare

Among other things, results above imply the existence of functorial injections

$$\mathbf{C}^\bullet[\mathcal{A}] \times \mathbf{C}^\bullet[\mathcal{B}] \hookrightarrow \mathbf{C}^\bullet[\mathcal{A} \circ \mathcal{B}], \quad \circ = \circ, \bullet, \odot, \triangleleft, \triangleright \text{ and } \diamond.$$

4.2.3. *Twist transformations and the coHom objects.* As an application of formulae above, given \mathcal{A} and \mathcal{B} in CA and a pair of 2nd admissible cocycles $\psi \in \mathfrak{Z}^2[\mathbf{A}_1]$ and $\varphi \in \mathfrak{Z}^2[\mathbf{B}_1]$, $\underline{\text{hom}}[\mathcal{B}_\varphi, \mathcal{A}_\psi] = \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_{\mathfrak{j}(\psi^\dagger, \varphi)}$. Then, for twisted coHom objects we have

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}_\varphi, \mathcal{A}_\psi] = \underline{\text{hom}}[\mathcal{B}_\varphi, \mathcal{A}_\psi]_\varsigma = \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_{\mathfrak{j}(\psi^\dagger, \varphi) \star \varsigma}.$$

On the other hand, from $\underline{\text{end}}[\mathcal{A}^{n|m}]$ and $\psi_{\mathfrak{h}}$ (see §3.1.1) we can compute the bialgebra $\underline{\text{end}}[\mathcal{A}_{\mathfrak{h}}^{n|m}]$ by making a twist transformation $\mathfrak{j}(\psi_{\mathfrak{h}}^\dagger, \psi_{\mathfrak{h}})$ (recall $\psi_{\mathfrak{h}}$ is a bicharacter). In general, as a direct consequence of **Theor. 16** for \triangleright :

Proposition 22. *If $\mathcal{A} \smile \mathcal{C}$ and $\mathcal{B} \smile \mathcal{D}$ through 2nd admissible cocycles, then $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \smile \underline{\text{hom}}[\mathcal{D}, \mathcal{C}]$. ■*

Now, we shall study twisted coHom objects under the perspective of twist transformation. From §3.1.4, we know that there exist admissible bicharacters (see Eqs. (3.14) and (3.15)), *ipso facto* 2nd admissible cochains, $\varsigma_{\mathcal{A}} \in \mathfrak{Z}^2[\mathbf{A}_1]$, $\varsigma_{\mathcal{B}} \in \mathfrak{Z}^2[\mathbf{B}_1]$ and $\varsigma_{\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]} \in \mathfrak{Z}^2[\mathbf{B}_1^* \otimes \mathbf{A}_1]$ such that

$$\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}] \simeq \mathcal{B}_\varsigma \triangleright \mathcal{A}_\varsigma = \underline{\text{hom}}[\mathcal{B}_\varsigma, \mathcal{A}_\varsigma] \simeq \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_\varsigma.$$

By direct calculations, it can be seen that $\varsigma_{\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]} = \mathfrak{j}(\varsigma_{\mathcal{B}}^\dagger, \varsigma_{\mathcal{A}})$. On the other hand, the product $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]_{\circ_\tau \mathcal{B}}$ appearing in twisted coevaluation map $\mathcal{A} \rightarrow \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]_{\circ_\tau \mathcal{B}}$, can be regarded as a twisting of the quantum space $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]_{\circ \mathcal{B}}$ by a cochain $\omega \in \mathfrak{Z}^2[\mathbf{B}_1^* \otimes \mathbf{A}_1 \otimes \mathbf{B}_1]$. The latter is an admissible anti-bicharacter defined by $\omega_{1,1}$ as (recall Eq. (3.13))

$$\omega(z_i^j \otimes z_r^s \otimes b_k \otimes b_l) = \phi_i^a \phi_r^b (z_a^c \otimes z_b^d) \rho_c^j \rho_d^s \otimes b_k \otimes b_l,$$

writing $z_i^j = b^j \otimes a_i$ and making usual identifications. Using (2.24) we find that $\omega = \mathfrak{j}(i\varsigma_{\mathcal{B}}^\dagger, i\varsigma_{\mathcal{A}}, \mathbb{I}^{\otimes 2})$. Since for all 2-cocycles ψ and φ ,

$$\mathfrak{j}(i\psi, i\varphi) = \mathfrak{j}\partial(\theta^{-1}, \chi^{-1}) = \partial(\mathfrak{j}(\theta, \chi)^{-1}) = \mathfrak{ij}(\psi, \varphi),$$

then $\omega = \mathfrak{ij}(\varsigma_{\mathcal{B}}^\dagger, \varsigma_{\mathcal{A}}, \mathbb{I}^{\otimes 2})$, and consequently

$$\begin{aligned} \underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]_{\circ_\tau \mathcal{B}} &= \left(\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_{\mathfrak{j}(\varsigma_{\mathcal{B}}^\dagger, \varsigma_{\mathcal{A}})} \circ \mathcal{B} \right)_\omega = ((\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \circ \mathcal{B})_{i\omega})_\omega \\ &= (\underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \circ \mathcal{B})_{i\omega \star \omega} = \underline{\text{hom}}[\mathcal{B}, \mathcal{A}] \circ \mathcal{B}. \end{aligned}$$

That is to say, the initiality of $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ follows from that of $\underline{\text{hom}}[\mathcal{B}, \mathcal{A}]$, because the related coevaluation maps are given by a same arrow. This way to see the construction of objects $\underline{\text{hom}}^\Upsilon[\mathcal{B}, \mathcal{A}]$ enable us to generalize it to more general twist transformations. In fact, instead of maps $\mathcal{A} \rightarrow \mathcal{H} \circ_\tau \mathcal{B}$ and twisting $\tau_{\mathcal{A}, \mathcal{B}}$, we can study a class of arrows $\mathcal{A} \rightarrow (\mathcal{H} \circ \mathcal{B})_\omega$, where ω defines an element of $\mathfrak{Z}^2[\mathbf{H}_1 \otimes \mathbf{B}_1]$. In terms of them, for each pair $\mathcal{A}, \mathcal{B} \in \text{CA}$, a full subcategory of $(\mathfrak{H}(\mathcal{A}) \downarrow \mathfrak{H}(\text{CA} \circ \mathcal{B}))$, being $\mathfrak{H} : \text{CA} \hookrightarrow \text{GrVct}$ the forgetful functor to graded vector spaces, can be defined as in [6]. If such a category, namely $\Omega^{\mathcal{A}, \mathcal{B}}$, is related to a cochain of the form $\omega_{\mathcal{A}, \mathcal{B}} = \mathfrak{ij}(\varsigma_{\mathcal{B}}^\dagger, \varsigma_{\mathcal{A}}, \mathbb{I}^{\otimes 2}) \in \mathfrak{Z}^2[\mathbf{B}_1^* \otimes \mathbf{A}_1 \otimes \mathbf{B}_1]$ (in a similar way that $\Upsilon^{\mathcal{A}, \mathcal{B}}$ is related to $\tau_{\mathcal{A}, \mathcal{B}}$),¹⁶ where $\varsigma_{\mathcal{A}}$ denotes a 2nd \mathcal{A} -admissible 2-cocycle, then it will have initial object

$$\underline{\text{hom}}^\Omega[\mathcal{B}, \mathcal{A}] = \underline{\text{hom}}[\mathcal{B}, \mathcal{A}]_{\mathfrak{j}(\varsigma_{\mathcal{B}}^\dagger, \varsigma_{\mathcal{A}})} = \mathcal{B}_\varsigma \triangleright \mathcal{A}_\varsigma.$$

Also, the disjoint union $\Omega = \bigvee_{\mathcal{A}, \mathcal{B}} \Omega^{\mathcal{A}, \mathcal{B}}$ has a semigroupoid structure, and there exists a related embedding $\Omega \hookrightarrow \text{CA}$ that preserves the corresponding (partial) products. Accordingly, the assignment $(\mathcal{B}, \mathcal{A}) \mapsto \underline{\text{hom}}^\Omega[\mathcal{B}, \mathcal{A}]$ defines an CA-cobased and (by duality) a QLS-based category for each collection of cocycles $\{\omega_{\mathcal{A}, \mathcal{B}}\}$. These results are developed in [7].

Finally, let us say the bialgebras $\underline{\text{end}}^\Omega[\mathcal{A}]$ are twisting of $\underline{\text{end}}[\mathcal{A}]$ also as bialgebras (i.e. in the Drinfeld's sense). In fact, the isomorphism given by Eq. (4.9) ensure the twisting $\mathfrak{j}(\varsigma^\dagger, \varsigma)$ on each bialgebra $\underline{\text{end}}[\mathcal{A}]$ is in the image of the map F , i.e. there exists $\chi \in \mathbf{G}^\bullet[\mathbf{A} \triangleright \mathbf{A}]$ such that $F_\chi = \mathfrak{j}(\varsigma^\dagger, \varsigma)$. Explicitly,

$$\chi_{\mathcal{A}}(z_{n_1}^{i_1} \dots z_{n_r}^{i_r} \otimes z_{m_1}^{k_1} \dots z_{m_s}^{k_s}) = (\varsigma_{\mathcal{A}})_{n_1 \dots n_r, m_1 \dots m_s}^{i_1 \dots i_r, k_1 \dots k_s}.$$

The fact that $\varsigma_{\mathcal{A}}$ is 2nd admissible is crucial for $\chi_{\mathcal{A}}$ to be well defined on $(\mathbf{A} \triangleright \mathbf{A})^{\otimes 2}$.

¹⁶To define $\omega_{\mathcal{A}, \mathcal{B}}$ in a diagram (φ, \mathcal{H}) , **Prop. 4** must be taken into account for the inclusions $\mathbf{H}_1 \otimes \mathbf{B}_1 \subset \mathbf{B}_1^* \otimes \mathbf{A}_1 \otimes \mathbf{B}_1$.

CONCLUSIONS

Motivated by the idea of twisted internal coHom objects, we have defined a non formal algebra deformation process over the category of quantum spaces. Such deformations, or twisting, are controlled by a cosimplicial multiplicative quasicomplex structure \mathbf{C}^\bullet in the category Grp_* of groups and unit preserving maps. Their counital 2-cocycles are the elements implementing the mentioned deformations. Also, given a quantum space \mathcal{A} , the cohomology classes of 2-cocycles ,through the so called \mathcal{A} -admissible cochains, defines the isomorphisms classes of twisting of \mathcal{A} .

We have shown that for each bialgebra in the category of quantum spaces, the corresponding \mathbf{C}^\bullet has the quasicomplex of bialgebra twist transformations as a subobject. More precisely, if \mathbf{V} is a coalgebra, then there exists a monic $\mathbf{G}_Z^\bullet[\mathcal{V}] \hookrightarrow \mathbf{C}^\bullet[\mathbf{V}]$ in the category of group quasicomplexes, where \mathcal{V} is a bialgebra generated by \mathbf{V} .

The twist transformations define a gauge equivalence between quantum spaces. For instance, every n -dimensional quantum plane $\mathcal{A}_h^{n|0}$ is gauge equivalent to the affine space $\mathcal{A}^{n|0}$. Also, the twisted coHom objects are equivalent, in this sense, to the untwisted ones.

The quasicomplexes $\mathbf{C}^\bullet[\mathbf{V}]$, $\mathbf{V} \in \text{Vct}_f$, generate a monoidal category with involution. We have shown that, for a particular class of twist transformations, the monoidal structure is compatible with products \circ , \bullet , \odot , \triangleleft , \triangleright and \diamond between quantum spaces, and the involution with the functor $!$, in the sense that these functors (acting on objects) ‘commute’ with the twisting process.

Now, it is natural to ask what happen if, instead of groups $\mathbf{C}^n[\mathbf{V}]$, we consider the algebras of degree zero n -homogeneous linear endomorphisms $\mathbf{E}^n[\mathbf{V}]$. Straightforwardly, the function $[n + 1] \mapsto \mathbf{E}^n[\mathbf{V}]$ together with Eqs. (2.6) and (2.10) define a cosimplicial object in Alg . We just have to repeat the proof of **Theor. 2**. Moreover, maps ∂_\pm given in Eq. (2.14), supply \mathbf{E}^\bullet with a structure of multiplicative cosimplicial parity quasicomplex in Alg_* (the category of algebras and unit preserving functions). From the algebra deformation process given at the beginning of §2, new twisting on quantum spaces can be constructed within the cohomological framework defined by $(\mathbf{E}^\bullet, \partial_+, \partial_-)$. These new transformations will be investigated elsewhere.

ACKNOWLEDGMENTS

The author thanks to CNEA, Argentina, for financial support, and thanks to H. Montani for useful discussions.

REFERENCES

- [1] A. Čap, H. Schichl, J. Vanžura, On twisted tensor products of algebras, *Comm. in Alg.* 23 (1995) 4701-4735.
- [2] J. Cornwell, A. Jacobs, Twisting 2-cocycles for the construction of new non-standard quantum groups, *J. Math. Phys.* 36 (1997) 5383-5401.
- [3] A. Davidov, Twisting of monoidal structures, pre-print math.QA/9703001.
- [4] V. G. Drinfeld, Quasi-Hopf algebras, *Leningrad Math J.* 2 (1990) 1419-1457.
- [5] M. Gerstenhaber, S.D. Shack, Algebra cohomology and deformation theory, in: *Deformation theory of algebras and structures, and applications*, Kluwer, 1988, pp. 11-264.
- [6] S. Grillo, H. Montani, Twisted internal coHom objects in the category of quantum spaces, pre-print math.QA/0112233.
- [7] S. Grillo, Generalized twisted coHom objects, math.QA/0204247.
- [8] G. Hochschild, On the cohomology groups of an associative algebra, *Annals of Math.* 46 (1945) 58-67.
- [9] L. Ionescu, On parity complexes and non-abelian cohomology, CT/980868.
- [10] S. Mac Lane, *Categories for the working mathematician*, Springer Verlag, 1971.
- [11] S. Majid, *Foundations of quantum group theory*, Cambridge University Press, 1995.
- [12] Yu. Manin, Some remarks on Koszul algebras and quantum groups, *Ann. Inst. Fourier* 33 (1987) 191-205.
Yu. Manin, *Quantum groups and non commutative geometry*, CRM, Université de Montréal, 1988.
- [13] Yu. Manin, *Topics in non commutative geometry*, Princeton University Press, 1991.
- [14] N.Yu. Reshetikhin, L.A. Takhtadjan, L.D. Faddeev, Quantization of Lie Groups and Lie Algebras, *Leningrad Math. J.* 1 (1990) 193-225.

Centro Atómico Bariloche and Instituto Balseiro, 8400-S. C. de Bariloche, Argentina
E-mail address: sergio@cabtep2.cnea.gov.ar